Patient Information

Montelukast (mon te loo' kast) Sodium Tablets

Read the Patient Information Leaflet that comes with montelukast sodium tablets before you start taking it and each time you get a refill. There may be new information. This leaflet does not take the place of talking with your healthcare provider about your medical condition or your treatment.

What is montelukast sodium tablet?

Montelukast sodium tablet is a prescription medicine that blocks substances in the body called leukotrienes. This
may help to improve symptoms of asthma and allergic rhinitis. Montelukast sodium tablets do not contain a
steroid.

Montelukast sodium tablets are used to:

1. Prevent asthma attacks and for the long-term treatment of asthma in adults and adolescents ages 15 years and

Do not take montelukast sodium tablets if you need relief right away for a sudden asthma attack. If you get an asthma attack, you should follow the instructions your healthcare provider gave you for treating asthma attacks.

- 2. Prevent exercise-induced asthma in people 15 years of age and older.
- 3. Help control the symptoms of allergic rhinitis (sneezing, stuffy nose, runny nose, itching of the nose). Montelukast sodium tablet is used to treat:
 - outdoor allergies that happen part of the year (seasonal allergic rhinitis) in adults and adolescents ages 15
 years and older, and
 - indoor allergies that happen all year (perennial allergic rhinitis) in adults and adolescents ages 15 years and older.

Who should not take montelukast sodium tablets?

Do not take montelukast sodium tablets if you are allergic to any of its ingredients.

See the end of this leaflet for a complete list of the ingredients in montelukast sodium tablets.

What should I tell my healthcare provider before taking montelukast sodium tablets?

Before taking montelukast sodium tablets, tell your healthcare provider if you:

- · are allergic to aspirin
- · have any other medical conditions
- are pregnant or plan to become pregnant. Talk to your doctor if you are pregnant or plan to become pregnant, as montelukast sodium tablets may not be right for you.
- are breast-feeding or plan to breast-feed. It is not known if montelukast passes into your breast milk. Talk to
 your healthcare provider about the best way to feed your baby while taking montelukast sodium tablets.

Tell your healthcare provider about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements. Some medicines may affect how montelukast sodium tablets works, or montelukast sodium tablets may affect how your other medicines work.

How should I take montelukast sodium tablets?

For anyone who takes montelukast sodium tablets:

- Take montelukast sodium tablets exactly as prescribed by your healthcare provider. Your healthcare provider will tell you how much montelukast sodium tablets to take, and when to take it.
- Do not stop taking montelukast sodium tablets or change when you take it without talking with your healthcare provider.
- · You can take montelukast sodium tablets with food or without food.
- If you or your child misses a dose of montelukast sodium tablets, just take the next dose at your regular time. Do not take 2 doses at the same time.
- If you take too much montelukast sodium tablets, call your healthcare provider or a Poison Control Center right away.

For adults and adolescents 15 years of age and older with asthma:

- Take montelukast sodium tablets 1 time each day, in the evening. Continue to take montelukast sodium tablets
 every day for as long as your healthcare provider prescribes it, even if you have no asthma symptoms.
- Tell your healthcare provider right away if your asthma symptoms get worse, or if you need to use your rescue inhaler medicine more often for asthma attacks.
- Do not take montelukast sodium tablets if you need relief right away from a sudden asthma attack. If you
 get an asthma attack, you should follow the instructions your healthcare provider gave you for treating asthma
 attacks.
- · Always have your rescue inhaler medicine with you for asthma attacks.
- Do not stop taking or lower the dose of your other asthma medicines unless your healthcare provider tells you to.

For patients 15 years of age and older for the prevention of exercise-induced asthma:

- · Take montelukast sodium tablets at least 2 hours before exercise.
- · Always have your rescue inhaler medicine with you for asthma attacks
- If you take montelukast sodium tablets every day for chronic asthma or allergic rhinitis, do not take another
 dose to prevent exercise-induced asthma. Talk to your healthcare provider about your treatment for exerciseinduced asthma.
- · Do not take 2 doses of montelukast sodium tablets within 24 hours (1 day).

For adults and adolescents 15 years of age and older with seasonal allergic rhinitis, or for adults and adolescents 15 years of age and older with perennial allergic rhinitis:

Take montelukast sodium tablets 1 time each day, at about the same time each day.

What is the dose of montelukast sodium tablets?

The dose of montelukast sodium tablets prescribed for your condition is based on age:

15 years and older: one 10 mg tablet.

What should I avoid while taking montelukast sodium tablets?

If you have asthma and aspirin makes your asthma symptoms worse, continue to avoid taking aspirin or other medicines called non-steroidal anti-inflammatory drugs (NSAIDs) while taking montelukast sodium tablets.

What are the possible side effects of montelukast sodium tablets?

Montelukast sodium tablets may cause serious side effects.

- · Behavior and mood-related changes. Tell your healthcare provider right away if you or your child have any of these symptoms while taking montelukast sodium tablets:
- · agitation including aggressive behavior or hostility
- · attention problems
- · bad or vivid dreams
- depression
- · disorientation (confusion)
- · feeling anxious
- hallucinations (seeing or hearing things that are not really there)
- · irritability
- · memory problems
- · restlessness
- sleep walking
- suicidal thoughts and actions (including suicide)
- · tremor
- · trouble sleeping
- · uncontrolled muscle movements
- Increase in certain white blood cells (eosinophils) and possible inflamed blood vessels throughout the body (systemic vasculitis). Rarely, this can happen in people with asthma who take montelukast sodium tablets. This sometimes happens in people who also take a steroid medicine by mouth that is being stopped or the dose is being lowered.

Tell your healthcare provider right away if you get one or more of these symptoms:

- · a feeling of pins and needles or numbness of arms or legs
- · a flu-like illness
- · severe inflammation (pain and swelling) of the sinuses (sinusitis)

The most common side effects with montelukast sodium tablets include:

- · upper respiratory infection
- · fever
- headache
- · sore throat
- cough
- · stomach pain
- diarrhea
- earache or ear infection

- flu
- · runny nose
- · sinus infection

Other side effects with montelukast sodium tablets include:

- increased bleeding tendency, low blood platelet count
- allergic reactions [including swelling of the face, lips, tongue, and/or throat (which may cause trouble breathing or swallowing), hives and itching]
- dizziness, drowsiness, pins and needles/numbness, seizures (convulsions or fits)
- palpitations
- nose bleed, stuffy nose, swelling (inflammation) of the lungs
- heartburn, indigestion, inflammation of the pancreas, nausea, stomach or intestinal upset, vomiting
- hepatitis
- bruising, rash, severe skin reactions (erythema multiforme, Stevens-Johnson syndrome/toxic epidermal necrolysis) that may occur without warning
- joint pain, muscle aches and muscle cramps
- bedwetting in children
- · tiredness, swelling

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of montelukast sodium tablets. For more information ask your healthcare provider or pharmacist.

Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store montelukast sodium tablets?

- Store montelukast sodium tablets at 20° to 25° C (68° to 77°F).
- Keep montelukast sodium tablets in the container it comes in.
- Keep montelukast sodium tablets in a dry place and away from light.

General Information about the safe and effective use of montelukast sodium tablets

Medicines are sometimes prescribed for purposes other than those mentioned in Patient Information Leaflets. Do not use montelukast sodium tabletsfor a condition for which it was not prescribed. Do not give montelukast sodium tablets to other people even if they have the same symptoms you have. It may harm them. **Keep montelukast and all medicines out of the reach of children.**

This leaflet summarizes information about montelukast sodium tablets. If you would like more information, talk to your healthcare provider. You can ask your pharmacist or healthcare provider for information about montelukast sodium tablets that is written for health professionals.

What are the ingredients in montelukast sodium tablets?

Active ingredient: montelukast sodium USP

Inactive ingredients: croscarmellose sodium, hydroxypropyl cellulose, lactose monohydrate, magnesium stearate and microcrystalline cellulose. The tablets are coated with opadry yellow which contains carnauba wax, hydroxypropyl cellulose, hypromellose, red iron oxide, titanium dioxide and yellow iron oxide.



Manufactured for: Camber Pharmaceuticals, Inc. Piscataway, NJ 08854

Ву: НЕТЕКОТМ

Hetero Labs Limited, Unit V, Polepally, Jadcherla, Mahabubnagar - 509 301, India.

Revised: 05/2017