

- **Stomach growths (fundic gland polyps).** People who take PPI medicines for a long time have an increased risk of developing a certain type of stomach growth called fundic gland polyps, especially after taking PPI medicines for more than 1 year.
- **Low magnesium levels in the body** can happen in people who have taken lansoprazole delayed-release capsules for at least 3 months. Tell your doctor right away if you have symptoms of low magnesium levels, including seizures, dizziness, irregular heartbeat, jitteriness, muscle aches or weakness, and spasms of hands or feet or voice.
- **Severe skin reactions.** Lansoprazole delayed-release capsules can cause rare but severe skin reactions that may affect any part of your body. These serious skin reactions may need to be treated in a hospital and may be life threatening:
 - Skin rash which may have blistering, peeling or bleeding on any part of your skin (including your lips, eyes, mouth, nose, genitals, hands and feet).
 - You may also have fever, chills, body aches, shortness of breath, or enlarged lymph nodes.

Stop taking lansoprazole delayed-release capsules and call your doctor right away. These symptoms may be the first sign of a severe skin reaction.

The most common side effects of lansoprazole delayed-release capsules include: diarrhea, stomach-ache (abdomen) pain, nausea and constipation. These are not all the possible side effects of lansoprazole delayed-release capsules. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store lansoprazole delayed-release capsules? Store lansoprazole delayed-release capsules at room temperature between 68°F to 77°F (20°C to 25°C).
Keep lansoprazole delayed-release capsules and all medicines out of the reach of children.

General information about the safe and effective use of lansoprazole delayed-release capsules. Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use lansoprazole delayed-release capsules for conditions for which it was not prescribed. Do not give lansoprazole delayed-release capsules to other people, even if they have the same symptoms that you have. It may harm them. You can ask your doctor or pharmacist for information about lansoprazole delayed-release capsules that is written for health professionals. For more information, call 1-866-495-1995.

What are the ingredients in lansoprazole delayed-release capsules?
Active ingredient: lansoprazole USP.
Inactive ingredients in lansoprazole delayed-release capsules: colloidal silicon dioxide, corn starch, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer dispersion, polysorbate 80, sucrose, sugar spheres (contains sucrose and starch (maltze)), talc, titanium dioxide and triethyl citrate. The hard gelatin capsule shell consists of gelatin, FD&C Blue No. 1, D&C Red No. 28, FD&C Red No. 40 and titanium dioxide. In addition 15 mg capsule contains FD&C Green No. 3. The imprinting ink contains polysorbate 80, propylene glycol, shellac and titanium dioxide.

Medication Guide available at
<http://camberpharma.com/medication-guides>

CAMBER
Pharmaceuticals, Inc.
Manufactured for:
Camber Pharmaceuticals, Inc.
Piscataway, NJ 08854
by: **HETERO™**
Hetero Labs Limited
Jeedimetla,
Hyderabad - 500 055, India

This Medication Guide has been approved by the U.S. Food and Drug Administration.
Revised: 09/2023

INSTRUCTIONS FOR USE

Lansoprazole (lan-SOH-pra zoh) Delayed-Release Capsules, USP

- Important:**
- Take lansoprazole delayed-release capsules before meals.
 - Do not crush or chew lansoprazole delayed-release capsules.
 - Lansoprazole delayed-release capsules should only be used with the foods and juices listed below.

Lansoprazole Delayed-Release Capsules
Taking lansoprazole delayed-release capsules with certain foods:
You can only use applesauce, ENSURE pudding, cottage cheese, yogurt or strained pears.

1. Open the capsule.
2. Sprinkle the granules on 1 tablespoon of applesauce, ENSURE pudding, cottage cheese, yogurt or strained pears.
3. Swallow right away.

Taking lansoprazole delayed-release capsules with certain juices:
You can only use apple juice, orange juice or tomato juice.

1. Open the capsule.
2. Sprinkle the granules into 60 mL (about ¼ cup) of apple juice, orange juice or tomato juice.
3. Stir.
4. Swallow right away.
5. To make sure that the entire dose is taken, add 1/2 cup or more of juice to the glass, stir and swallow right away.

Giving lansoprazole delayed-release capsules through a nasogastric tube (NG tube) size 16 French or larger:

- You can only use apple juice.**
1. Place 40 mL of apple juice into a clean container.
 2. Open the capsule and empty the granules into the container of apple juice.
 3. Use a catheter-tip syringe to draw up the apple juice and granule mixture.
 4. Gently mix the catheter-tip syringe to keep the granules from settling.
 5. Attach the catheter-tip syringe to the NG tube.
 6. Give the mixture right away through the NG tube that goes into the stomach. Do not save the apple juice and granule mixture for later use.
 7. Refill the catheter-tip syringe with 40 mL of apple juice and mix gently. Flush the NG tube with apple juice.

How should I store lansoprazole delayed-release capsules?

- Store lansoprazole delayed-release capsules at room temperature between 68°F to 77°F (20°C to 25°C).

Keep lansoprazole delayed-release capsules and all medicines out of the reach of children.

This Instruction for Use has been approved by the U.S. Food and Drug Administration. All brand names listed are the registered trademarks of their respective owners and are not trademarks of Hetero Labs Limited.

CAMBER
Pharmaceuticals, Inc.
Manufactured for:
Camber Pharmaceuticals, Inc.
Piscataway, NJ 08854
by: **HETERO™**
Hetero Labs Limited
Jeedimetla,
Hyderabad - 500 055, India

Revised: 09/2023

Non-erectile GERD Improvement in Overall GERD Symptoms*	71.2% (42/59)†
Erosive Esophagitis Improvement in Overall GERD Symptoms*	78.2% (18/23)†
Healing Rate†	85.2% (21/22)†

* Symptoms assessed by patient diary (parents/caregivers as necessary).
† No data available for five patients.
‡ Data from one healed patient was excluded from this analysis due to timing of final endoscopy.

In these 87 adolescent patients, increases in serum gastrin levels were similar to those observed in adult studies, median fasting serum gastrin levels increased 42% from 45 pM at baseline to 64 pM. [Interquartile range (25th to 75th percentile) of 44 to 88 pM] at the final visit. (Normal serum gastrin levels are 25 to 111 pM.)

The safety of lansoprazole delayed-release capsules has been assessed in these 87 adolescent patients. Of the 87 adolescent patients with GERD, 6% (5/87) took lansoprazole for less than six weeks, 52% (45/87) for six to 10 weeks, and 1% (1/87) for greater than 10 weeks. The most frequently reported (at least 3%) treatment-related adverse reactions in these patients were headache (7%), abdominal pain (5%), nausea (3%) and dizziness (3%). Treatment-related dizziness, reported in this prescribing information as occurring in less than 1% of adult patients, was reported in this study by three adolescent patients with non-erectile GERD, who had dizziness associated with other reactions (such as migraine, dyspnea, and vomiting).

Juvenile Animal Toxicity Data
Heart Valve Thickening
In two oral toxicity studies, thickening of the mitral heart valve occurred in juvenile rats treated with lansoprazole. Heart valve thickening was observed primarily with oral dosing initiated on postnatal Day 1 (age equivalent to neonatal humans) and postnatal Day 14 (human age equivalent of approximately one year) at doses of 250 mg/kg/day and higher (at postnatal Day 7 and postnatal Day 14, respectively 6.2 times and 4.2 times the daily pediatric dose of 15 mg in pediatric patients age one to 11 years weighing 30 kg or less, based on AUC). The treatment durations associated with heart valve thickening ranged from 5 days to 8 weeks. The findings observed or trended towards reversibility after a 4-week drug-free recovery period.

The incidence of heart valve thickening after initiation of dosing on postnatal Day 21 (human equivalent of approximately 11 years) was not observed in juvenile rats. In a group given 500 mg/kg/day for 4 or 8 weeks (approximately 5.2 times the daily pediatric dose of 15 mg in pediatric patients age one to 11 years weighing 30 kg or less, based on AUC). Based on exposure margins, the risk of heart valve injury does not appear to be relevant to patients of one year of age and older.

Bone Changes
In an eight-week oral toxicity study in juvenile rats with dosing initiated on postnatal Day 7, doses equal to or greater than 100 mg/kg/day (2.5 times the daily pediatric dose of 15 mg in children age one to 11 years weighing 30 kg or less, based on AUC) produced dose-dependent, with impairment of weight gain observed as early as postnatal Day 10 (age equivalent to neonatal humans). At the end of treatment, the signs of impaired growth at 100 mg/kg/day and higher included reductions in body weight (14 to 44% compared to controls), absolute weight gains of approximately 10 to 15% compared to controls, and reductions in the length of multiple organs, femur weight, femur length, and crown-rump length. Femoral growth plate thickness, which is only AUC-related, was reduced by approximately 10 to 15% compared to control growth persisted through the end of the four-week recovery period. Longer term data were not collected.

6.5 Genetic Use
Of the total number of patients (n=21,486) in clinical studies of lansoprazole, 16% of patients were aged 65 years and over, while 4% were 75 years and over. No overall differences in safety or effectiveness were observed between these patients and younger patients and other reported clinical experience has not identified significant differences in responses between children and younger patients given greater sensitivity of some older populations cannot be ruled out [see Clinical Pharmacology (12.3)].


6.6 Hepatic Impairment
In patients with varying degrees of chronic hepatic impairment the exposure to lansoprazole was increased compared to healthy subjects with normal hepatic function [see Clinical Pharmacology (12.3)]. No dosage adjustment for lansoprazole is necessary for patients with mild (Child-Pugh Class A) or moderate (Child-Pugh Class B) hepatic impairment. The recommended dosage is 15 mg orally daily for patients with severe hepatic impairment (Child-Pugh Class C) [see Dosage and Administration (2.3)].

10 OVERDOSEAGE
Lansoprazole is not removed from the circulation by hemodialysis. In one reported overdose, a patient consumed 600 mg of lansoprazole capsules in a single episode. The patient was treated with 5,000 mg of i.v. naltrexone 1 hour (approximately 1.3 times the 30 mg human dose based on body surface area (BSA)) and received approximately 61.7 times the 30 mg human dose based on BSA) did not produce clinically or any clinical effects (FDA Green No. 3).

In the event of over-exposure, treatment should be symptomatic and supportive.

If over-exposure occurs, call your poison control center at 1-800-222-1222 for current information regarding assessment of poisoning or over-exposure.

11 DESCRIPTION
The active ingredient in lansoprazole delayed-release capsules, USP, is lansoprazole, a substituted benzimidazole, 2-[(5S)-6-methoxy-3,4-dihydro-2H-1,2,4-benzodiazepin-5-ylidene]-1H-benzimidazole, a compound that inhibits gastric acid secretion. The empirical formula is C₁₆H₁₄F₃N₂O₃ with a molecular weight of 359.36. Lansoprazole has the following structure:



Lansoprazole, USP is a white to brownish-white powder which melts with decomposition at approximately 168°C. Lansoprazole is freely soluble in dimethylformamide and practically insoluble in water.

The rate of degradation of the compound in aqueous solution increases with decreasing pH.

Lansoprazole is supplied in delayed-release capsules for oral administration.

Lansoprazole delayed-release capsules are available in two dosage strengths: 15 mg and 30 mg of lansoprazole delayed-release capsules, USP per capsule. Each delayed-release capsule contains enteric-coated granules consisting of 15 mg or 30 mg of lansoprazole (active ingredient) and the following inactive ingredients: colloidal silicon dioxide, corn starch, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer dispersion, polysorbate 80, sucrose, sugar spheres (contains sucrose and starch (maltze)), talc, titanium dioxide and triethyl citrate. The hard gelatin capsule shell consists of gelatin, FD&C Blue No. 1, D&C Red No. 28, FD&C Red No. 40 and titanium dioxide. In addition 15 mg capsule contains FD&C Green No. 3.

The imprinting ink contains polysorbate 80, propylene glycol, shellac and titanium dioxide.

12 CLINICAL PHARMACOLOGY
12.1 Mechanism of Action
Lansoprazole belongs to a class of antisercretory compounds, the substituted benzimidazoles, that suppress gastric acid secretion by specific inhibition of the H⁺-K⁺-ATPase enzyme system at the apical surface of the gastric parietal cell. Because this enzyme is responsible for the acid pump (proton) pump within the parietal cell, lansoprazole has been characterized as a gastric acid pump inhibitor. In addition to the inhibition of gastric acid secretion, this effect also leads to inhibition of both basal and stimulated gastric acid and pepsin secretion of the stimulus. Lansoprazole does not exhibit anticholinergic or histamine type-2 antagonist activity.

12.2 Pharmacodynamics
Antisecretory Activity
After oral administration, lansoprazole was shown to significantly decrease the basal acid output and significantly increase the basal gastric pH and lower gastric volume. Oral lansoprazole was greater than three and greater than four. Lansoprazole also significantly reduced meal-stimulated gastric output and secretory volume, as well as gastric volume (Table 7).

In patients with hypersecretion of acid, lansoprazole significantly reduced basal and pentagastrin-stimulated gastric acid secretion. Lansoprazole inhibited the normal increase in secretion volume, acidity and acid output induced by insulin.

The intragastric pH results of a five day, pharmacodynamic, crossover study of 15 and 30 mg of once daily lansoprazole are presented in Table 8.

Parameter	Lansoprazole				
	Baseline Value	15 mg twice daily	30 mg twice daily	30 mg	30 mg
Mean 24 Hour pH	2.1	2.7	4.0	3.8	4.9
Mean Nighttime pH	1.9	2.4	3.0	2.6	3.8
% Time Gastric pH>3	18	33	59*	51*	72*
% Time Gastric pH>4	12	22	48*	41*	66*

NOTE: An intragastric pH of greater than four reflects a reduction in gastric acidity by 98%.
(p<0.05) vs baseline only.
(p<0.05) vs baseline and lansoprazole 15 mg.

After the initial dose of this study, increased gastric pH was seen in one to two hours with 30 mg of lansoprazole and two to three hours with 15 mg of lansoprazole. After multiple daily dosing, increased gastric pH was seen within the first four postdosing with 30 mg of lansoprazole and within one to two hours postdosing with 15 mg of lansoprazole.

Acid suppression may enhance the effect of antimicrobials in eradicating *Helicobacter pylori* (H. pylori). The percentage of time gastric pH was elevated (above 4) was evaluated in a crossover study of lansoprazole given daily, twice and three times daily (Table 7).

Parameter	Lansoprazole				
	30 mg daily	15 mg twice daily	30 mg twice daily	30 mg	30 mg three times
% Time Gastric pH>5	43	47	59*	57*	71*
% Time Gastric pH>6	20	23	28	45*	47*

(p<0.05) vs lansoprazole 30 mg daily.
(p<0.05) vs lansoprazole 30 mg daily, 15 and 30 mg twice daily.

The inhibition of gastric acid secretion by lansoprazole is gradually returned to normal over one to four days after multiple doses. There was no indication of rebound gastric acidity.

Endothelinoformin-like (EC₂) Cell Effects
During limited exposure of rats with up to 150 mg/kg/day of lansoprazole doses seven days per week, marked hypersecretion was observed followed by EC₂ cell proliferation and increased levels of endothelinoformin in female rats. Gastric tumor species in the fundus of the stomach from approximately 150 patients treated continuously with lansoprazole for at least one year did not show evidence of EC₂ cell effects similar to those seen in rat studies. Longer term data are needed to rule out the possibility of an increased risk of the development of gastric tumors in patients receiving long-term therapy with lansoprazole [see Carcinogenicity (12.1)].

Other Gastric Effects in Humans
Lansoprazole did not significantly affect mucosal blood flow in the fundus of the stomach. Due to the normal physiological effect of gastric acid on the pyloric sphincter, there was a decrease of about 17% in blood flow in the antrum, pylorus, and duodenal bulb was seen. Lansoprazole significantly slowed the gastric emptying of digestible solids. Lansoprazole increased serum pepsinogen levels and decreased pepsin activity under basal conditions and in response to meal stimulation or insulin injection. As with other agents that elevate intragastric pH, increase in gastric pH was associated with increased in nitrite-reducing bacteria and elevation of nitrite concentration in gastric juice in patients with gastric ulcer. No significant increase in nitrosamine concentrations was observed.

Serum Gastrin Effects
In over 2,100 patients, median fasting serum gastrin levels increased 50 to 100% from baseline but remained within normal range after treatment with 15 to 60 mg of oral lansoprazole. These elevations reached a plateau within two months of therapy and returned to pretreatment levels within four weeks after discontinuation of therapy.

Increased gastric acid secretion enhances cell hyperplasia and increased serum CgA levels. The increased CgA levels may cause false positive results in diagnostic investigations for neuroendocrine tumors [see Warnings and Precautions (5.3)].

Endocrine Effects
Human studies for up to one year have not detected any clinically significant effects on the endocrine system. Hormones studied include testosterone, luteinizing hormone (LH), follicle stimulating hormone (FSH), sex hormone binding globulin (SHBG), dehydroepiandrosterone sulfate (DHEA-S), prolactin, cortisol, estradiol, insulin, aldosterone, parathormone, growth stimulating hormone (GSH), triiodothyronine (T₃), thyroxine (T₄), and somatotrophic hormone (STH). Lansoprazole in oral doses of 15 to 60 mg for up to one year had no clinically significant effect on sexual function. In addition, lansoprazole in oral doses of 15 to 60 mg for two to eight weeks had no clinically significant effect on thyroid function. In 24-month carcinogenicity studies in Sprague-Dawley rats with daily lansoprazole dosages up to 150 mg/kg, proliferative changes in the Leydig cells of the testes, including benign neoplasms, were increased compared to control rats.

Other Effects
No systemic effects of lansoprazole on the central nervous system, lymphoid, hematopoietic, renal, hepatic, cardiovascular, or respiratory systems have been observed in humans. Among 56 patients who had extensive baseline eye evaluations, no visual toxicity was observed after lansoprazole treatment (up to 180 mg/day) for up to 58 to 60 weeks. In addition, 15 to 60 mg of oral lansoprazole, the peak plasma concentrations (C_{max}) of lansoprazole and the area under the plasma concentration curves (AUCs) of lansoprazole were approximately proportionally to the administered dose. Lansoprazole does not accumulate and is not metabolized by blocking the proton pump (H⁺-K⁺-ATPase enzyme system) at the apical surface of the gastric parietal cell. The two active species are not present in the systemic circulation. The plasma elimination half-life of lansoprazole is less than two hours while the acid inhibitory effect lasts more than 24 hours. Therefore, the plasma elimination half-life of lansoprazole does not reflect its duration of suppression of gastric acid secretion.

Excitation: Following single-dose oral administration of lansoprazole, virtually no unchanged lansoprazole was excreted in the urine. In one study, after a single oral dose of 40 mg lansoprazole, approximately one-third of the administered radiolabel was excreted in the urine and one-third was recovered in the feces. This implies a significant biliary excretion of the lansoprazole metabolites.

Specific Populations
Pediatric Patients:
Over 17 years of age
The pharmacokinetics of lansoprazole were studied in pediatric patients with GERD aged one to 11 years and 12 to 17 years in two separate clinical studies. In children aged one to 11 years, lansoprazole was dosed 15 mg daily for subjects weighing 30 kg and 30 mg daily for subjects weighing greater than 30 kg. Mean C_{max} and AUC values observed on Day 5 of dosing were similar between the two dose groups and were not affected by weight or age within each weight-adjusted dose group used in the study. In adolescent subjects aged 12 to 17 years, subjects were randomized to receive lansoprazole at 15 or 30 mg daily. Mean C_{max} and AUC values of lansoprazole were not affected by body weight or age, and nearly dose-proportional increases in mean C_{max} and AUC values were observed between the two dose groups in the study. Overall, lansoprazole pharmacokinetics in pediatric patients aged one to 17 years were similar to those observed in healthy adult subjects.

Geriatric Patients:
The clearance of lansoprazole is decreased in the elderly, with elimination half-life increased approximately 50 to 100%. Because the mean half-life in the elderly remains between 1.3 to 2.6 hours, repeated once daily dosing does not result in accumulation of lansoprazole. Peak plasma levels were not increased in the elderly [see Use in Specific Populations (8.3)].

Male and Female Patients:
In a study comparing 12 male and six female human patients who received lansoprazole, no sex-related differences were found in pharmacokinetics and nitroglycerin pH results.

Racial or Ethnic Groups:
The pooled mean pharmacokinetic parameters of lansoprazole from twelve US studies (N=513) were compared to the mean pharmacokinetic parameters from two Asian studies (N=20). The mean AUCs of lansoprazole in Asian subjects were approximately twice those seen in pooled US data; however, the inter-individual variability was high. The C_{max} values were comparable.

Renal Impairment:
In patients with severe renal impairment, plasma protein binding decreased by 1 to 1.5% after administration of 60 mg of lansoprazole. Patients with renal impairment had a shortened elimination half-life and decreased total AUC. Free and bound AUC for free lansoprazole in plasma, however, was not related to the degree of renal impairment; and the C_{max} and T_{max} (time to reach the maximal concentration) were not affected. Because the C_{max} and T_{max} data from subjects with normal renal function. Therefore, the pharmacokinetics of lansoprazole do not clinically differ in patients with mild, moderate or severe renal impairment compared to healthy subjects with normal function.

Patients with Hepatic Impairment:
In patients with mild (Child-Pugh Class A) or moderate (Child-Pugh Class B) hepatic impairment, there was an approximate 3-fold increase in mean AUC compared to healthy subjects with normal hepatic function following multiple oral doses of 30 mg lansoprazole for seven days. The corresponding mean plasma half-life of lansoprazole was prolonged from 1.5 to four hours (Child-Pugh A) or five to nine hours (Child-Pugh B).

In patients with compensated and decompensated cirrhosis, there was an approximate 10-fold increase in mean AUC, respectively, compared to healthy subjects with normal hepatic function following a single oral dose of 30 mg lansoprazole [see Dosage and Administration (2.3) and Use in Specific Populations (8.6)].

Drug Interaction Studies
Effect of Lansoprazole on Other Drugs
Cytochrome P450 Interactions:
Lansoprazole is metabolized through the cytochrome P450 system, specifically through the CYP2C19 and CYP2C19 isozymes. Studies have shown that lansoprazole does not have clinically significant interactions with other drugs metabolized by the cytochrome P450 system, such as diazepam, aspirin, indomethacin, ibuprofen, phenytoin, propofol, prednisone, diazepam, or theophylline in healthy subjects. These compounds are metabolized through various cytochrome P450 isozymes including CYP1A2, CYP2C18, CYP2C8, CYP2C9, and CYP3A4.

When lansoprazole was administered concomitantly with theophylline (CYP1A2, CYP2C19, and CYP2C9), there was no clinically significant interaction. Because of the similar magnitude and the direction of the effect on theophylline clearance, this interaction is unlikely to be of clinical concern [see Drug Interactions (7)].

Metabolism and Excretion:
In an open-label, single-arm, eight day, pharmacokinetic study of 28 adult rheumatoid arthritis patients who required the chronic use of 7.5 to 15 mg of methotrexate given weekly (12 mg/m² by oral administration), the pharmacokinetics of lansoprazole 30 mg daily had no effect on the pharmacokinetics of methotrexate and 7-hydroxymethotrexate. While this study was not designed to assess the safety of lansoprazole in patients receiving methotrexate, reactions were noted. However, this study was conducted with low doses of methotrexate. A drug interaction study with high doses of methotrexate has not been conducted [see Warnings and Precautions (7.1)].

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been shown to have no clinically significant interaction with anticancer drugs.

Anticancer:
Lansoprazole has also been