

Oxcarbazepine Tablets USP, for oral use
(ox'' kar baz' e peen)

- nausea
- tiredness (lack of energy)
- headache
- confusion
- more frequent or more severe seizures

Your healthcare provider may do blood tests to check your sodium levels during your treatment with oxcarbazepine tablets.

- swelling of your face, eyes, lips, or tongue
- trouble swallowing or breathing
- a skin rash
- hives
- fever, swollen glands, or sore throat that do not go away or come and go
- painful sores in the mouth or around your eyes
- yellowing of your skin or eyes
- unusual bruising or bleeding
- severe fatigue or weakness
- severe muscle pain
- frequent infections or infections that do not go away

- thoughts about suicide or dying
- attempts to commit suicide
- new or worse depression
- new or worse anxiety
- feeling agitated or restless
- panic attacks
- trouble sleeping (insomnia)
- new or worse irritability
- acting aggressive, being angry, or violent
- acting on dangerous impulses
- an extreme increase in activity and talking (mania)
- other unusual changes in behavior or mood

- Stopping oxcarbazepine tablets suddenly can cause serious problems.
- Stopping a seizure medicine suddenly in a patient who has epilepsy may cause seizures that will not stop (status epilepticus).


- alone or with other medicines to treat partial-onset seizures in adults
- alone to treat partial-onset seizures in children 4 years and older
- with other medicines to treat partial-onset seizures in children 2 years and older

- have or have had suicidal thoughts or actions, depression or mood problems
- have liver problems
- have kidney problems

- are allergic to carbamazepine. Many people who are allergic to carbamazepine are also allergic to oxcarbazepine tablets.
- use birth control medicine. Oxcarbazepine tablets may cause your birth control medicine to be less effective. Talk to your healthcare provider about the best birth control method to use.

- are pregnant or plan to become pregnant. Oxcarbazepine tablets may harm your unborn baby. Tell your healthcare provider right away if you become pregnant while taking oxcarbazepine tablets. You and your healthcare provider will decide if you should take oxcarbazepine tablets while you are pregnant.

If you become pregnant while taking oxcarbazepine tablets, talk to your healthcare provider about registering with the North American Antiepileptic Drug (NAAED) Pregnancy Registry. The purpose of this registry is to collect information about the safety of antiepileptic medicine during pregnancy. You can enroll

Artwork information			
Customer	Camber	Market	USA
Dimensions (mm)	210 x 350 mm	Non Printing Colors	 Die cut
Pharma Code No.	NA		
Printing Colours	Black		
Others: NA			

in this registry by calling 1-888-233-2334.

- are breastfeeding or plan to breastfeed. Oxcarbazepine passes into breast milk. Talk with your healthcare provider about the best way to feed your baby if you take oxcarbazepine tablets.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking oxcarbazepine tablets with certain other medicines may cause side effects or affect how well they work. Do not start or stop other medicines without talking to your healthcare provider.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

- How should I take oxcarbazepine tablets?**
- Do not stop taking oxcarbazepine tablets without talking to your healthcare provider. Stopping oxcarbazepine tablets suddenly can cause serious problems, including seizures that will not stop (status epilepticus).
 - Take oxcarbazepine tablets exactly as prescribed. Your healthcare provider may change your dose. Your healthcare provider will tell you how many oxcarbazepine tablets to take.
 - Take oxcarbazepine tablets 2 times a day.
 - Take oxcarbazepine tablets with or without food.
 - If you take too much oxcarbazepine, call your healthcare provider right away.

- What should I avoid while taking oxcarbazepine tablets?**
- Do not drive or operate machinery until you know how oxcarbazepine tablets affect you. Oxcarbazepine tablets may slow your thinking and motor skills.
 - Do not drink alcohol or take other drugs that make you sleepy or dizzy while taking oxcarbazepine tablets until you talk to your healthcare provider. Oxcarbazepine tablets taken with alcohol or drugs that cause sleepiness or dizziness may make your sleepiness or dizziness worse.

- What are the possible side effects of oxcarbazepine tablets?**
- See “What is the most important information I should know about oxcarbazepine tablets?”
- Oxcarbazepine tablets may cause other serious side effects, including:**
- trouble concentrating
 - problems with your speech and language
 - feeling confused
 - feeling sleepy and tired
 - trouble with walking and coordination
 - seizures that can happen more often or become worse, especially in children

- Get medical help right away if you have any of the symptoms listed above or listed in “What is the most important information I should know about oxcarbazepine tablets?”**
- The most common side effects of oxcarbazepine tablets include:**
- dizziness
 - sleepiness
 - double vision
 - tiredness
 - nausea
 - vomiting
 - problems with vision
 - trembling
 - problems with walking and coordination (unsteadiness)
 - rash

These are not all the possible side effects of oxcarbazepine tablets. Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

- How should I store oxcarbazepine tablets?**
- Store oxcarbazepine film-coated tablets at room temperature between 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C to 30°C (59°F to 86°F). Keep oxcarbazepine tablets film-coated tablets dry.
- Keep oxcarbazepine tablets and all medicines out of the reach of children.**

General Information about the safe and effective use of oxcarbazepine tablets.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use oxcarbazepine tablets for a condition for which it was not prescribed. Do not give oxcarbazepine tablets to other people, even if they have the same symptoms that you have. They may harm them.


You can ask your pharmacist or healthcare provider for information about oxcarbazepine tablets that is written for health professionals.

What are the ingredients in oxcarbazepine tablets?

Active ingredient: oxcarbazepine

Inactive ingredients: colloidal silicon dioxide, crospovidone, hypromellose, magnesium stearate, microcrystalline cellulose, black iron oxide, iron oxide yellow, iron oxide red, polyethylene glycol, polyvinyl alcohol, talc, and titanium dioxide.

Medication Guide available at <http://camberpharma.com/medication-guides>



Manufactured for:
Camber Pharmaceuticals, Inc.
Piscataway, NJ 08854.

By: Annora Pharma Pvt. Ltd.
Sangareddy - 502313, Telangana, India.

For more information, call Annora Pharma Private Limited at 1-866-495-1995.