

4/2023

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use LACOSAMIDE INJECTION safely and effectively. See full prescribing tion for LACOSAMIDE INJECTION LACOSAMIDE injection, for intravenous use, CV

-- INDICATIONS AND USAGE-

Initial U.S. Approval: 2008 -- RECENT MAJOR CHANGES Dosage and Administration (2.2)

Lacosamide is indicated for: Treatment of partial-onset seizures in patients 17 years of age and older (1.1)

-- DOSAGE AND ADMINISTRATION

Adults (17 years and older) Initial dosage for monotherapy for the treatment of partial-onset seizures is $100\,\mathrm{mg}$ twice daily (2.1)

Initial dosage for adjunctive therapy for the treatment of partial-onset seizures is 50 mg twice daily (2.1) $\label{lem:maximum recommended dosage for monotherapy and adjunctive therapy is 200\,mg \ twice \ daily \ (2.1)$ Increase do sage based on clinical response and tolerability, no more frequently than once per week (2.1)

Injection: for intravenous use only when oral administration is temporarily not feasible; the reco times daily over 15 to 60 minutes; obtaining ECG before initiation is recommended in certain patients (2.7, 5.3) Dose adjustment is recommended for severe renal impairment (2.4, 12.3)

adjustment is recommended for mild or moderate hepatic impairment; use in patients with severe hepatic impairment is not reco ...DOSAGE FORMS AND STRENGTHS... 200 mg/20 mL single-dose vial for intravenous use (3)

-- CONTRAINDICATIONS

None (4)

Monitor natients for suicidal behavior and ideation (5.1)

Cardiac Rhythm and Conduction Abnormalities: Obtaining ECG before beginning and after titration to steady-state maintenance is recommended patients with underlying proarrhythmic conditions or on concomitant medications that affect cardiac conduction: closely monitor these

-- WARNINGS AND PRECAUTIONS

Lacosamide on Johannia (Lacosamide Caralla) withdrawn to minimize the potential of increased seizure frequency (5.5)

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Multi-Organ Hypersensitivity: Discontinue if no alternate etiology (5.6)

·····ADVERSE REACTIONS····

somnolence (6.1) Monotherapy: Most common adverse reactions are similar to those seen in adjunctive therapy studies (6.1)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide Pediatric use information is approved for UCB, Inc.'s VIMPAT® (lacosamide) injection. However, due to UCB, Inc.'s marketing exclusivity rights, this

FULL PRESCRIBING INFORMATION: CONTENTS*

- INDICATIONS AND USAGE
- DOSAGE AND ADMINISTRATION
 - Dosage Information Alternate Initial Dosage Information
- Converting From a Single Antiepileptic (AED) to Lacosamide Injection Monotherapy for the Treatment of Partial-Onset Seizure: Dosage Information for Patients with Renal Impairmen
- 2.5 Dosage Information for Patients with Hepatic Impairment Preparation and Administration Information for Lacosamide Injection
- 2.8 Discontinuation of Lacosamide Injection
- 3 DOSAGE FORMS AND STRENGTHS
- CONTRAINDICATIONS WARNINGS AND PRECAUTIONS
- Suicidal Behavior and Ideation
- Cardiac Rhythm and Conduction Abnormalities Withdrawal of Antiepileptic Drugs (AEDs)
- ADVERSE REACTIONS 6.2 Postmarketing Experience
- DRUG INTERACTIONS
- 7.1 Strong CYP3A4 or CYP2C9 Inhibitors
 7.2 Concomitant Medications that Affect Cardiac Conduction USE IN SPECIFIC POPULATIONS
- 8.1 Pregnancy 8.2 Lactation

drug product is not labeled with that pediatric information

FULL PRESCRIBING INFORMATION 1 INDICATIONS AND USAGE

Lacosamide injection is indicated for the treatment of partial-onset seizures in patients 17 years of age and older Pediatric use information is approved for UCB, Inc.'s VIMPAT® (lacosamide) injection. However, due to UCB, Inc.'s marketing exclusivity rights, this

5.6 Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Multi-Organ Hypersensitivity

2 DOSAGE AND ADMINISTRATION

2.1 Dosage Informat The recommended dosage for monotherapy and adjunctive therapy for partial-onset seizures in 17 years of age and older is included in Table 1. Dosage should be increased based on clinical response and tolerability, no more frequently than once per week. Titration increments should not exceed those

Table 1: Recommended Dosages for Partial-Onset Seizures (Monotherapy or Adjunctive Therapy) in Patients 17 years of Age and Older Age and Body Weight **Initial Dosage Titration Regimen** Maintenance Dosage Adults (17 years and older) Monotherapy **: 100 mg twice Monotherapy **: 150 mg to 200 mg Increase by 50 mg twice daily daily (200 mg per day) twice daily (300 mg to 400 mg per day Adjunctive Therapy: 50 mg twice Adjunctive Therapy: 100 mg to 200 mg daily (100 mg per day) twice daily (200 mg to 400 mg per day) *when not specified, the dosage is the same for monotherapy for partial-onset seizures and adjunctive therapy for partial-onset seizures

*Monotherapy for partial-onset seizures only

In adjunctive clinical trials in adult patients with partial-onset seizures, a dosage higher than 200 mg twice daily ($400 \,\mathrm{mg}$ per day) was not more effective and was associated with a substantially higher rate of adverse reactions (see Adverse Reactions (6.1) and Clinical Studies (14.2)). Lacosamide Injection Dosage

Lacosamide injection may be used when oral administration is temporarily not reasible *[see Dosage and Administration (2.0) and Warnings (Precautions (5.3)]*. Lacosamide injection can be administered intravenously to adult patients with the same dosing regimens described for oral dosing. The clinical study experience of intravenous lacosamide is limited to 5 days of consecutive treatment Pediatric use information is approved for UCB, Inc.'s VIMPAT® (lacosamide) injection. However, due to UCB, Inc.'s marketing exclusivity rights, this

drug product is not labeled with that pediatric information 2.2 Alternate Initial Dosage Information to Achieve the Maintenance Dosage in a Shorter Timeframe herapy and adjunctive therapy for partial-onset seizures in patients 17 years of age and older, an alternate initial dosing regimen for week 1

(e.g., including a loading dose and/or a higher initial dosage) may be administered in patients for whom achieving the recommended maintenance dosage in a shorter timeframe is clinically indicated (see Table 2). The alternate initial dosage regimen should be continued for one week. Lacosamide injection may then be titrated based on clinical response and tolerability, no more frequently than once per week, if needed. The loading dose should be administered with medical supervision because of the possibility of increased incidence of adverse reactions, including central nervous system (CNS) and cardiovascular adverse reactions (see Warnings and Precautions (5.2, 5.3), Adverse Reactions (6.1), and Clinical Pharmacology (12.3). Titration crements should not exceed those shown in Table 2. Table 2: Alternate Initial Dosing Regimen to Achieve the Maintenance Dosage in a Shorter Timeframe if Clinically Indicated*

Age and Body Weight	Alternate Initial Dosage	Titration Regimen	Maintenance Dosage
Adults (17 years and older)	Single loading dose: 200 mg 12 hours later initiate: 100 mg twice daily (200 mg per day)	Increase by 50 mg twice daily (100 mg per day) at weekly intervals, if needed	Monotherapy**: 150 mg to 200 mg twice daily (300 mg to 400 mg per day) Adjunctive Therapy: 100 mg to 200 mg twice daily (200 mg to 400 mg per day)

*when not specified, the dosage is the same for monotherapy for partial-onset seizures and adjunctive therapy for partial-onset seizures
**Monotherapy for partial-onset seizures only

2.3 Converting From a Single Antienilentic (AED) to Lacosamide Injection Monotherany for the Treatment of Partial-Onset Seizures For patients who are already on a single AED and will convert to lacosamide injection monotherapy, withdrawal of the concomitant AED should until the therapeutic dosage of lacosamide injection is achieved and has been administered for at least 3 days. A gradual withdrawal of the concomitan AED over at least 6 weeks is recommended.

2.4 Dosage Information for Patients with Renal Impairment For patients with mild to moderate renal impairment, no dosage adjustment is necessary. For patients with severe renal impairment [creatinine clearance (CL_{ca}) less than 30 mL/min as estimated by the Cockcroft-Gault equation for adults; CLCR less than 30 ml/min/1.73m² as estimated by the Schwartz equation for pediatric patients] or end-stage renal disease, a reduction of 25% of the maximum dosage is

In all patients with renal impairment, dose initiation and titration should be based on clinical response and tolerability **Hemodialysis** Lacosamide injection is effectively removed from plasma by hemodialysis. Following a 4-hour hemodialysis treatment, dosage supplementation of up to

50% should be considered. Concomitant Strong CYP3A4 or CYP2C9 Inhibitors Dose reduction may be necessary in patients with renal impairment who are taking strong inhibitors of CYP3A4 and CYP2C9 [see Drug Interactions (7.1), Use in Specific Populations (8.6), and Clinical Pharmacology (12.3)].

2.5 Dosage Information for Patients with Hepatic Impairmen For patients with mild or moderate hepatic impairment, a reduction of 25% of the maximum dosage is recommended. The dose initiation and titration should be based on clinical response and tolerability performed with caution in patients with hepatic impairment.

Concomitant Strong CYP3A4 and CYP2C9 Inhibitors se reduction may be necessary in patients with hepatic impairment who are taking strong inhibitors of CYP3A4 and CYP2C9 (see Drug Interaction.

(7.1), Use in Specific Populations (8.7), and Clinical Pharmacology (12.3)]. 2.7 Preparation and Administration Information for Lacosamide Injection

Lacosamide injection can be administered intravenously without further dilution or may be mixed with diluents listed below. The diluted solution should

Sodium Chloride Injection 0.9% (w/v) Dextrose Injection 5% (w/v)

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Product with particulate matter or discoloration should not be used.

d infusion duration is 30 to 60 minutes; however, infusions as rapid as 15 minutes can be administered in adults if required *(see Adverse* Reactions (6.1) and Clinical Pharmacology (12.3)]. Infusion durations less than 30 minutes are generally not recommended in pediatric patients /see

Intravenous infusion of lacosamide may cause bradycardia, AV blocks, and ventricular tachyarrhythmia (see Warnings and Precautions (5.3/). Obtaining an ECG before beginning lacosamide injection and after lacosamide injection is titrated to steady-state maintenance dose is recomm with underlying prograthythmic conditions or on concomitant medications that affect cardiac conduction (see Drug Interactions (7.2)).

tion should not be stored for more than 4 hours at room temperature. Any unused portion of lacosamide injection should be discarded.2.8 Discontinuation of Lacosamide injection

When discontinuing lacosamide injection, a gradual withdrawal over at least 1 week is recommended (see Warnings and Precautions (5.5)). 3 DOSAGE FORMS AND STRENGTHS

Lacosamide Injection, USP 200 mg/20 mL; clear, colorless sterile solution in single-dose vials 4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS 5.1 Suicidal Behavior and Ideation

Antiepileptic drugs (AEDs), including lacosamide, increase the risk of suicidal thoughts or behavior in patients taking these drugs for any indication Patients treated with any AED for any indication should be monitored for the emergence or worsening of depression, suicidal thoughts or behavior, and/or

Pooled analyses of 199 placebo-controlled clinical trials (mono- and adjunctive therapy) of 11 different AEDs showed that patients randomized to one of the AEDs had approximately twice the risk (adjusted Relative Risk 1.8, 95% Cl:1.2, 2.7) of suicidal thinking or behavior compared to patients rando to placebo. In these trials, which had a median treatment duration of 12 weeks, the estimated incidence of suicidal behavior or ideation among 27,863 AFD-treated nationts was 0.43%, compared to 0.24% among 16.029 placeho-treated nationts, representing an increase of approximately one case of suicidal thinking or behavior for every 530 patients treated. There were four suicides in drug-treated patients in the trials and none in placebo-treated patients, but the number of events is too small to allow any conclusion about drug effect on suicide. The increased risk of suicidal thoughts or behavior with AEDs was observed as early as one week after starting treatment with AEDs and persisted for

the duration of treatment assessed. Because most trials included in the analysis did not extend beyond 24 weeks, the risk of suicidal thoughts or vior beyond 24 weeks could not be assessed.

The risk of suicidal thoughts or behavior was generally consistent among drugs in the data analyzed. The finding of increased risk with AEDs of varying mechanisms of action and across a range of indications suggests that the risk applies to all AEDs used for any indication. The risk did not vary substantially by age (5 to 100 years) in the clinical trials analyzed.

Table 3 shows absolute and relative risk by indication for all evaluated AEDs

Indication	Placebo Patients with Events Per 1000 Patients	Drug Patients with Events Per 1000 Patients	Relative Risk: Incidence of Events in Drug Patients/ Incidence in Placebo Patients	Risk Difference: Additional Drug Patients with Events Per 1000 Patients
Epilepsy	1.0	3.4	3.5	2.4
Psychiatric	5.7	8.5	1.5	2.9
Other	1.0	1.8	1.9	0.9
Total	2.4	4.3	1.8	1.9

Anyone considering prescribing lacosamide or any other AED must balance this risk with the risk of untreated illness. Epilepsy and many other illnesses ileptics are prescribed are themselves associated with morbidity and mortality and an increased risk of suicidal thoughts and behavior Should suicidal thoughts and behavior emerge during treatment, the prescriber needs to consider whether the emergence of these symptoms in any given patient may be related to the illness being treated 5.2 Dizziness and Ataxia

Lacosamide may cause dizziness and ataxia in adult and pediatric patients. In adult patients with partial-onset seizures taking 1 to 3 concomitant AEDs, dizziness was experienced by 25% of patients randomized to the recommended doses (200 to 400 mg/day) of lacosamide (compared with 8% of placebo patients) and was the adverse reaction most frequently leading to discontinuation (3%). Ataxia was experienced by 6% of patients randomized to the recommended doses (200 to 400 mg/day) of lacosamide (compared to 2% of placebo patients). The onset of dizziness and ataxia was most comr observed during titration. There was a substantial increase in these adverse reactions at doses higher than 400 mg/day *[see Adverse Reactions (6.1)]*. If a loading dose is clinically indicated, administer with medical supervision because of the possibility of increased incidence of adverse reactions, including CNS adverse reactions such as dizziness and ataxia.

5.3 Cardiac Rhythm and Conduction Ab $\underline{PR\,Interval\,Prolongation,\,Atrioven tricular\,Block,\,and\,Ventricular\,Tachyarrhythmia}$

Size: 350 x 730 mm

Colour: Black

Dose-dependent prolongations in PR interval with lacosamide have been observed in clinical studies in adult patients and in healthy volunteers (see Dose-upperioring projection in mice var with autosamine lance been observed been desired in a final clinical studies in adult patients and in meantly outnities (See Clinical Pharmacology (12.2)), in adjunctive clinical trials in adult patients with partial-onset seizures, asymptomatic first-degree atrioventricular (AV) block was observed as an adverse reaction in 0.4% (4/944) of patients randomized to receive lacosamide and 0% (0/364) of patients randomized to receive placebo. One case of profound bradycardia was observed in a patient during a 15-minute infusion of 150 mg lacosamide. When lacosamide is given with other drugs that prolong the PR interval, further PR prolongation is possible. In the postmarketing setting, there have been reports of cardiac arrhythmias in patients treated with lacosamide, including bradycardia, AV block, and

ventricular tachyarrhythmia, which have rarely resulted in asystole, cardiac arrest, and death. Most, although not all, cases have occurred in patients with underlying proarrhythmic conditions, or in those taking concomitant medications that affect cardiac conduction or prolong the PR interval. These events have occurred with both oral and intravenous routes of administration and at prescribed doses as well as in the setting of overdose [see Overdosage (10)). In all patients for whom a loading dose is clinically indicated, administer the loading dose with medical supervision because of the $possibility\ of\ increased\ incidence\ of\ adverse\ reactions,\ including\ cardiov ascular\ adverse\ reactions.$ Lacosamide should be used with caution in patients with underlying proarrhythmic conditions such as known cardiac conduction problems (e.g., marked

first-degree AV block, second-degree or higher AV block and sick sinus syndrome without pacemaker), severe cardiac disease (such as myccardial ischemia or heart failure, or structural heart disease), and cardiac sodium channelopathies (e.g., Brugada Syndrome). Lacosamide should also be used with caution in patients on concomitant medications that affect cardiac conduction, including sodium channel blockers, beta-blockers, calcium channe ekers, potassium channel blockers, and medications that prolong the PR interval [see Drug Interactions (7.2)]. In such patients, obtaining an ECG before beginning lacosamide, and after lacosamide is titrated to steady-state maintenance dose, is recommended. In addition, these patients should be closely monitored if they are administered lacosamide through the intravenous route (see Adverse Reactions (6.1) and Drug Interactions (7.2)). Atrial Fibrillation and Atrial Flutter

ional trials of lacosamide in adult patients with partial-onset seizures there were no cases of atrial fibrillation or flutter. Both atrial fibrillation and atrial flutter have been reported in open label partial-onset seizure trials and in postmarketing experience. In adult patients with diabetic neuropathy, for which lacosamide is not indicated, 0.5% of patients treated with lacosamide experienced an adverse reaction of atrial fibrillation or atrial flutter, compared to 0% of placebo-treated patients. Lacosamide administration may predispose to atrial arrhythmias (atrial fibrillation or flutter), especially in patients with diabetic neuropathy and/or cardiovascular disease.

syncope compared to placebo. In the short-term controlled trials in adult patients with diabetic neuropathy, for which lacosamide is not indicated, 1.2% of patients who were treated with lacosamide reported an adverse reaction of syncope or loss of consciousness, compared with 0% of placebo-treated patients with diabetic neuropathy. Most of the cases of syncope were observed in patients receiving doses above 400 mg/day. The cause of syncope was not determined in most cases. However, several were associated with either changes in orthostatic blood pressure, atrial flutter/fibrillation (and associated tachycardia, or bradycardia. Sees of syncope have also been observed in open-label clinical partial-onset seizure studies in adult and pediatric patients. These cases were associated with a history of risk factors for cardiac disease and the use of drugs that slow AV conduction. 5.5 Withdrawal of Antienilentic Drugs (AFDs)

As with all AEDs, lacosamide should be withdrawn gradually (over a minimum of 1 week) to minimize the potential of increased seizure frequency in natients with seizure disorders 5.6 Drug Reaction with Fosinophilia and Systemic Symptoms (DRESS)/Multi-Organ Hypersensitivity

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS), also known as multi-organ hypersensitivity, has been reported in patients taking antiepileptic drugs, including lacosamide. Some of these events have been fatal or life-threatening. DRESS typically, although not exclusively, presents with fever, rash, lymphadenopshy and/or facial swelling, in association with other organ system involvement, such as hepatitis, hepatris, benatologic abnormalities, myocarditis, or myositis, sometimes resembling an acute viral infection. Eosinophilia is often present. This disorder is variable in its expression, and other organ systems not noted here may be involved. It is important to note that early manifestations of hypersensitivity (e.g., fever, lymphadenopathy) may be present even though rash is not evident. If such signs or symptoms are present, the patient should be evaluated immediately.

- Lacosamide may cause syncope (5.4)
- Adjunctive therapy: Most common adverse reactions in adults (≥ 10% and greater than placebo) are diplopia, headache, dizziness, nausea, and
- To report SUSPECTED ADVERSE REACTIONS, contact Aspiro Pharma Limited at 1-866-495-1995 or FDA at 1-800-FDA-1088 or
- ... USE IN SPECIFIC POPULATIONS Pregnancy: Based on animal data, may cause fetal harm (8.1)

Renal Impairment

- DRUG ABUSE AND DEPENDENCE Controlled Substance
- 9.3 Dependence
- 10 OVERDOSAGE 11 DESCRIPTION

8.4 Pediatric Use

- 12 CLINICAL PHARMACOLOGY
- 12.2 Pharmacodynamics 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY
- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 14 CLINICAL STUDIES 1 Monotherapy in Patients with Partial-Onset Seizures
- 14.2 Adjunctive Therapy in Patients with Partial-Onset Seizures 16 HOW SUPPLIED/STORAGE AND HANDLING
- 16.2 Storage and Handling 17 PATIENT COUNSELING INFORMATION

Lacosamide should be discontinued if an alternative etiology for the signs or symptoms cannot be established 6 ADVERSE REACTIONS

The following serious adverse reactions are described below and elsewhere in the labeling: Suicidal Behavior and Ideation (see Warnings and Precautions (5.1))

- Dizziness and Ataxia (see Warnings and Precautions (5.2))
- Cardiac Rhythm and Conduction Abnormalities [see Warnings and Precautions (5.3)] Syncope (see Warnings and Precautions (5.4)) Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Multiorgan Hypersensitivity Reactions/see Warnings and Precautions (5.6)/
- 6.1 Clinical Trials Experience Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly

compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice taking 1 to 3 concomitant anti-epileptic drugs. Elevations of ALT to ≥ 3x ULN occurred in 0.7% (7/935) of lacosamide patients and 0% (0/356) of ebo patients. One case of hepatitis with transaminases > 20x ULN occurred in one healthy subject 10 days after lacosamide treat completion, along with nephritis (proteinuria and urine casts). Serologic studies were negative for viral hepatitis. Transaminases returned to

normal within one month without specific treatment. At the time of this event, bilirubin was normal. The hepatitis/nephritis was interpreted as a

The following is a list of adverse reactions reported by patients treated with lacosamide in all clinical trials in adult patients, including controlled trials and long-term open-label extension trials. Adverse reactions addressed in other tables or sections are not listed here

Blood and lymphatic system disorders: neutropenia, anemia Cardiac disorders: palpitations

Gastrointestinal disorders: constipation, dyspepsia, dry mouth, oral hypoaesthesi

General disorders and administration site conditions: irritability, pyrexia, feeling drunk Injury, poisoning, and procedural complications: fall Musculoskeletal and connective tissue disorders: muscle spasms

Nervous system disorders: paresthesia, cognitive disorder, hypoaesthesia, dysarthria, disturbance in attention, cerebellar syndrome

Table 5 gives the incidence of adverse reactions that occurred in \geq 5% of adult patients in any lacosamide dosing group

Adult Patients (17 Years and Older Adverse reactions with intravenous administration to adult patients with partial-onset seizures generally were similar to those that occurred with the oral formulation, although intravenous administration was associated with local adverse reactions such as injection site pain or discomfort (2.5%), irritation (1%), and erythema (0.5%). One case of profound bradycardia (26 bpm: BP 100/60 mmHg) occurred in a patient during a 15 minute infusion ol 150 mg lacosamide. This patient was on a beta-blocker. Infusion was discontinued and the patient experienced a rapid recovery The safety of a 15-minute loading dose administration of lacosamide Injection 200 mg to 400 mg followed by oral administration of lacosamide given

twice daily at the same total daily dose as the initial intravenous infusion was assessed in an open-label study in adult patients with partial-onsel seizures. Patients had to have been maintained on a stable dose regimen of 1 to 2 marketed antiepileptics for at least 28 days prior to treatment assignment. Treatment groups were as follows: Single dose of intravenous lacosamide Injection 200 mg followed by oral lacosamide 200 mg/day (100 mg every 12 hours Single dose of intravenous lacosamide Injection 300 mg followed by oral lacosamide 300 mg/day (150 mg every 12 hours Single dose of intravenous lacosamide Injection 400 mg followed by oral lacosamide 400 mg/day (200 mg every 12 hours)

	Lacosamide 200 mg	Lacosamide 300 mg	Lacosamide 400 mg	Lacosamide Total N = 100
	N = 25	N=50	N=25	%
Adverse Reaction	%	%	%	
Eye disorders				
Diplopia	4	6	20	9
Blurred Vision	0	4	12	5
Gastrointestinal disorders				
Nausea	0	16	24	14
Dry mouth	0	6	12	6
Vomiting	0	4	12	5
Oral Paresthesia	4	4	8	5
Oral Hypoesthesia	0	6	8	5
Diarrhea	0	8	0	4
General disorders/administration site				
conditions				
Fatigue	0	18	12	12
Gait disturbance	8	2	0	3
Chest pain	0	0	12	3
Nervous system disorders				
Dizziness	20	46	60	43
Somnolence	0	34	36	26
Headache	8	4	16	8
Paresthesia	8	6	4	6
Tremor	0	6	4	4
Abnormal Coordination	0	6	0	3
Skin & subcutaneous tissue disorders				
Pruritus	0	6	4	4
Hyperhidrosis	0	0	8	2

Adverse reactions that occurred with infusion of lacosamide 200 mg over 15-minutes followed by lacosamide 100 mg administered orally twice per day were similar in frequency to those that occurred in 3-month adjunctive therapy controlled trials. Considering the difference in period of observations (1 week vs. 3 months), the incidence of CNS adverse reactions, such as dizziness, somnolence, and paresthesia may be higher with 15-minute administration of lacosamide Injection than with administration over a 30- to 60-minute period.

Pediatric use information is approved for UCB, Inc.'s VIMPAT® (lacosamide) injection. However, due to UCB, Inc.'s marketing exclusivity rights, thi: drug product is not labeled with that pediatric infor

The following adverse reactions have been identified during post-approval use of lacosamide. Because these reactions are reported voluntarily from a

opulation of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposur Blood and lymphatic system disorders: Agranulocytosis Psychiatric disorders: Aggression, agitation, hallucination, insomnia, psychotic disorder

Skin and subcutaneous tissue disorders: Angioedema, rash, urticaria, Stevens-Johnson syndrome, toxic epidermal necrolysis.

DRUG INTERACTIONS

7.1 Strong CYP3A4 or CYP2C9 Inhibitors Patients with renal or hepatic impairment who are taking strong inhibitors of CYP3A4 and CYP2C9 may have a significant increase in exposure to lacosamide. Dose reduction may be necessary in these patients. 7.2 Concomitant Medications that Affect Cardiac Conduction

Lacosamide should be used with caution in patients on concomitant medications that affect cardiac conduction (sodium channel blockers, beta-blockers, calcium channel blockers, potassium channel blockers) including those that prolong PR interval (including sodium channel blocking AEDs), because of a risk of AV block, bradycardia, or ventricular tachyarrhythmia. In such patients, obtaining an ECG before beginning lacosamide, and after lacosamide is titrated to steady-state, is recommended. In addition, these patients should be closely monitored if they are administered lacosamide through the intravenous route [see Warnings and Precautions (5.3)].

8 USE IN SPECIFIC POPULATIONS 8.1 Pregnancy Pregnancy Exposure Registry

Clinical Considerations

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to antiepileptic drugs (AEDs), such as lacosamide, during pregnancy. Encourage women who are taking lacosamide during pregnancy to enroll in the North American Antiepileptic Drug (NAAED) pregnancy registry by calling 1-888-233-2334 or visiting http://www.aedpregnancyregistry.org/.

Available data from the North American Antiepileptic Drug (NAAED) pregnancy registry, a prospective cohort study, case reports, and a case series with lacosamide use in pregnant women are insufficient to identify a drug associated risk of major birth defects, miscarriage or other adverse maternal or fetal outcomes. Lacosamide produced developmental toxicity (increased embryofetal and perinatal mortality, growth deficit) in rats following administration during pregnancy. Developmental neurotoxicity was observed in rats following administration during a period of postnatal development corresponding to the third trimester of human pregnancy. These effects were observed at doses associated with clinically relevant plasma exposu The background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Oral administration of lacosamide to pregnant rats (20, 75, or 200 mg/kg/day) and rabbits (6.25, 12.5, or 25 mg/kg/day) during the period of organogenesis did not produce any effects on the incidences of fetal structural abnormalities. However, the maximum doses evaluated were limited by ernal toxicity in both species and embryofetal death in rats. These doses were associated with maternal plasma lacosamide exposures (AUC) approximately 2 and 1 times (rat and rabbit, respectively) that in humans at the maximum recommended human dose (MRHD) of 400 mg/day. In two studies in which lacosamide (25, 70, or 200 mg/kg/day and 50, 100, or 200 mg/kg/day) was orally administered to rats throughout pregnancy are lactation, increased perinatal mortality and decreased body weights in the offspring were observed at the highest dose tested. The no-effect dose for pre- and postnatal developmental toxicity in rats (70 mg/kg/day) was associated with a maternal plasma lacosamide AUC similar to that in humans at the MRHD. Oral administration of lacosamide (30, 90, or 180 mg/kg/day) to rats during the neonatal and juvenile periods of development resulted in decreased brain

weights and long-term neurobehavioral changes (altered open field performance, deficits in learning and memory). The early postnatal period in rats is

generally thought to correspond to late pregnancy in humans in terms of brain development. The no-effect dose for developmental neurotoxicity in rats

In Vitro Data differentiation and control of axonal outgrowth. Potential adverse effects on CNS development related to this activity cannot be ruled out

8.2 Lactation Risk Summary Data from published literature indicate that lacosamide is present in human milk. There are reports of increased sleepiness in breastfed infants exposed to lacosamide (see Clinical Considerations). There is no information on the effects of lacosamide on milk production The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for lacosamide and any potentia

Monitor infants exposed to lacosamide through breastmilk for excess sedation

was associated with a plasma lacosamide AUC less than that in humans at the MRHD.

Safety and effectiveness in pediatric patients below 1 month of age, have not been established.

Lacosamide has been shown in vitro to interfere with the activity of collapsin response mediator protein-2 (CRMP-2), a protein involved in neuronal differentiation and control of axonal outgrowth. Potential clied to leated adverse effects on CNS development cannot be ruled out. Administration of lacosamide to rats during the neonatal and juvenile periods of postnatal development (approximately equivalent to neonatal through adolescent development in humans) resulted in decreased brain weights and long-term neurobehavioral changes (altered open field performance, deficits in learning urotoxicity in rats was associated with a plasma lacosamide exposure (AUC) less than that humans at the maximum recommended human dose of 400 mg/day. Pediatric use information is approved for UCB, Inc.'s VIMPAT® (lacosamide) injection. However, due to UCB, Inc.'s marketing exclusivity rights, this drug product is not labeled with that pediatric information

There were insufficient numbers of elderly patients enrolled in partial-onset seizure trials (n=18) to adequately determine whether they respond No lacosamide dose adjustment based on age is necessary. In elderly patients, dose titration should be performed with caution, usually starting at the lower end of the dosing range, reflecting the greater frequency of decrea sed hepatic function, decreased r abnormalities, and polypharmacy (see Dosage and Administration (2.1, 2.4, 2.5) and Clinical Pharmacology (12,3)]. 8.6 Renal Impairment

No dose adjustment is necessary in patients with mild to moderate renal impairment (CL , s \geq 30 mL/min). In patients with severe renal impairment (CL , s < 30 mL/min as estimated by the Cockcroft-Gault equation for adults) and in those with end-stage renal disease, a reduction of 25% of the maximum dosage is recommended [see Dosage and Administration (2.4) and Clinical Pharmacology (12.3)]. In all patients with renal impairment, dose initiation and titration should be based on clinical response and tolerability Lacosamide is effectively removed from plasma by hemodialysis. Dosage supplementation of up to 50% following hemodialysis should be considered 9.7 Henstie Impairmen

For adult and pediatric patients with mild to moderate hepatic impairment, a reduction of 25% of the maximum dosage is recommended. Patients with mild to moderate hepatic impairment should be observed closely for adverse reactions, and dose initiation and titration should be based on clinical response and tolerability [see Dosage and Administration (2.5), Clinical Pharmacology (12.3)]. The pharmacokinetics of lacosamide has not been evaluated in severe hepatic impairment. Lacosamide use is not recommended in patients with severe

9 DRUG ABUSE AND DEPENDENCE 9.1 Controlled Substance Lacosamide is a Schedule V controlled substance

Abuse is the intentional, non-therapeutic use of a drug, even once, for its desirable psychological or physiological effects. In a human abuse potentia

study, single doses of 200 mg (equal to the maximum single dosage) and 800 mg lacosamide (equal to twice the recommended daily maintenance dosage) produced euphoria-type subjective responses that differentiated statistically from placebo; at 800 mg, these euphoria-type responses were statistically indistinguishable from those produced by alprazolam, a Schedule IV drug. The duration of the euphoria-type responses following lacosamide was less than that following alprazolam. A high rate of euphoria was also reported as an adverse event in the human abuse potential study following single doses of 800 mg lacosamide (15% [5/34]) compared to placebo (0%) and in two pharmacokinetic studies following single and multiple doses of 300 to 800 mg lacosamide (ranging from 8% [2/33] to 25% [3/12]) compared to placebo (0%). However, the rate of euphoria reported as an adverse event in the lacosamide program at therapeutic doses was less than 1%. 9.3 Dependence Physical dependence is a state that develops as a result of physiological adaptation in response to repeated drug use, manifested by withdrawal signs and symptoms after abrupt discontinuation or a significant dose reduction of a drug. Abrupt termination of lacosamide in clinical trials with diabetic

neuropathic pain patients produced no signs or symptoms that are associated with a withdrawal syndrome indicative of physical dependence. However

you get a refill. There may be new information. This Medication Guide describes important safety information about lacosamide injection. This information does not take the place of talking to your healthcare provider about your medical condition or

MEDICATION GUIDE

Lacosamide (la koe' sa mide)

injection, USP

for intravenous use, CV

Read this Medication Guide before you start taking lacosamide injection and each time

What is the most important information I should know about lacosamide

Do not stop taking lacosamide injection without first talking to your healthcare provider. Stopping lacosamide injection suddenly can cause serious problems. Stopping seizure medicine suddenly in a patient who has epilepsy can cause seizures that will not stop (status epilepticus).

Lacosamide injection can cause serious side effects, including:

especially if they are new, worse, or worry you:

new or worse anxiety

panic attacks

feeling agitated or restless

thoughts, or feelings.

1. Like other antiepileptic drugs, lacosamide injection may cause suicidal thoughts or actions in a very small number of people, about 1 in 500.

Call a healthcare provider right away if you have any of these symptoms,

 thoughts about suicide or dying
 trouble sleeping (insomnia) attempt to commit suicide new or worse irritability new or worse depression

acting aggressive, being angry, or

 acting on dangerous impulses an extreme increase in activity and talking (mania)

 other unusual changes in behavior or mood

How can I watch for early symptoms of suicidal thoughts and actions? · Pay attention to any changes, especially sudden changes, in mood, behaviors,

 Keep all follow-up visits with your healthcare provider as scheduled. • Call your healthcare provider between visits as needed, especially if you are worried about symptoms.

• Suicidal thoughts or actions can be caused by things other than medicines. If

you have suicidal thoughts or actions, your healthcare provider may check for 2. Lacosamide injection may cause you to feel dizzy, have double vision, feel sleepy, or have problems with coordination and walking. Do not drive, operate heavy

machinery, or do other dangerous activities until you know how lacosamide injection affects you. 3. Lacosamide injection may cause you to have an irregular heartbeat or may cause you to faint. In rare cases, cardiac arrest has been reported. Call your healthcare

provider right away if you: have a fast, slow, or pounding heartbeat or feel your heart skip a beat

have shortness of breath

 have chest pain feel lightheaded

4. Lacosamide is a federally controlled substance (CV) because it can be abused or lead to drug dependence. Keep your lacosamide injection in a safe place, to protect it

• to treat partial-onset seizures in people 17 years of age and older.

fainted or if you feel like you are going to faint

harm them. Selling or giving away this medicine is against the law. What is lacosamide injection? Lacosamide injection is a prescription medicine used:

If you have fainted or feel like you are going to faint you should lay down with your

from theft. Never give your lacosamide injection to anyone else, because it may

It is not known if lacosamide injection is safe and effective for partial-onset seizures in children under 1 month of age.

What should I tell my healthcare provider before taking lacosamide injection? Before you take lacosamide injection, tell your healthcare provider about all of your medical conditions, including if you:

• have or have had depression, mood problems or suicidal thoughts or behavior. have heart problems. have kidney problems. have liver problems.

• have abused prescription medicines, street drugs or alcohol in the past.

while taking lacosamide injection. You and your healthcare provider will decide if you should take lacosamide injection while you are pregnant. o If you become pregnant while taking lacosamide injection, talk to your healthcare provider about registering with the North American Antiepileptic Drug Pregnancy Registry. You can enroll in this registry by calling 1-888-233-2334. The purpose of this registry is to collect information about the safety of

• are pregnant or plan to become pregnant. It is not known if lacosamide can harm

your unborn baby. Tell your healthcare provider right away if you become pregnant

antiepileptic medicine during pregnancy. are breastfeeding or plan to breastfeed. Lacosamide passes into breast milk.

o Breastfeeding during treatment with lacosamide injection may cause your baby to have more sleepiness than normal. If this happens, contact your baby's

o Talk to your healthcare provider about the best way to feed your baby if you take lacosamide injection.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking lacosamide injection with certain other medicines may cause side effects or affect how well they work. Do not start or stop other medicines without talking to your healthcare provider. Know the medicines you take. Keep a list of them and show it to

your healthcare provider and pharmacist each time you get a new medicine.

seizures that will not stop (status epilepticus).

How should I take lacosamide injection? • Take lacosamide injection exactly as your healthcare provider tells you.

 Your healthcare provider will tell you how much lacosamide to take and when to Your healthcare provider may change your dose if needed. • Do not stop lacosamide injection without first talking to a healthcare provider. Stopping lacosamide injection suddenly in a patient who has epilepsy can cause

 Lacosamide injection may be taken with or without food. • If you take too much lacosamide, call your healthcare provider or local Poison Control Center right away. What should I avoid while taking lacosamide injection?

how lacosamide injection affects you. Lacosamide injection may cause you to feel dizzy, have double vision, feel sleepy, or have problems with coordination and walking.

Do not drive, operate heavy machinery, or do other dangerous activities until you know

What are the possible side effects of lacosamide injection? • See "What is the most important information I should know about

lacosamide injection?" Lacosamide injection may cause other serious side effects including: • A serious allergic reaction that may affect your skin or other parts of your body such as your liver or blood cells. Call your healthcare provider right away if

o swelling of the legs o a skin rash, hives o fever or swollen glands that do not o yellowing of the skin or whites of

you have:

o tiredness (fatigue)

headache

the eyes o shortness of breath o dark urine

The most common side effects of lacosamide injection include: double vision

 dizziness These are not all of the possible side effects of lacosamide injection. For more information ask your healthcare provider or pharmacist. Tell your healthcare provider about any side effect that bothers you or that does not go away. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store lacosamide injection? • Store lacosamide injection at room temperature between 68°F to 77°F (20°C to

25°C). Do not freeze lacosamide injection.

Keep lacosamide injection and all medicines out of the reach of children.

General Information about the safe and effective use of lacosamide injection. Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use lacosamide injection for a condition for which it was not prescribed. Do not give lacosamide injection to other people, even if they have the same symptoms that you have. It may harm them.

This Medication Guide summarizes the most important information about lacosamide

injection. If you would like more information, talk with your healthcare provider. You can

ask your pharmacist or healthcare provider for information about lacosamide injection

that is written for health professionals. What are the ingredients in lacosamide injection?

Active ingredient: lacosamide Injection inactive ingredients: sodium chloride (7.60 mg/mL), water for injection,

hydrochloric acid (for pH adjustment).

Pharma Code: Front Side: 2220 Back Side: 2220

Pediatric use information is approved for UCB, Inc.'s VIMPAT® (lacosamide) injection. However, due to UCB, Inc.'s marketing exclusivity rights, this drug product is not labeled with that pediatric information

Medication Guide available at http://camberpharma.com/medicationguides

Manufactured for: Camber Pharmaceuticals, Inc., Piscataway, NJ 08854

Manufactured by: **Aspiro Aspiro Pharma Limited** Survey No. 321, Biotech Park, Phase – III Karkapatla Village, Markook (Mandal)

Siddipet, Telangana-502281, India.

For more information, call 1-866-495-1995

This Medication Guide has been approved by the U.S. Food and Drug Administration

Revised: 05/2023

psychological dependence cannot be excluded due to the ability of lacosamide to produce euphoria-type adverse events in humans 10 OVERDOSAGE

Events reported after an intake of more than 800 mg (twice the maximum recommended daily dosage) of lacosamide include dizziness, nausea, and seizures (generalized tonic clonic seizures, status epilepticus). Cardiac conduction disorders, confusion, decreased level of consciousness, cardiogenic shock, cardiac arrest, and coma have also been observed. Fatalities have occurred following lacosamide overdoses of several grams. There is no specific antidote for overdose with lacosamide. Standard decontamination procedures should be followed. General supportive care of the patient is indicated including monitoring of vital signs and observation of the clinical status of patient. A Certified Poison Control Center should be contacted for up to date information on the management of overdose with lacosamide.

indicated based on the patient's clinical state or in patients with significant renal impairment.

11 DESCRIPTION The chemical name of lacosamide, the single (R)-enantiomer, is (R)-2-acetamido-N-benzyl-3-methoxypropionamide (IUPAC). Lacosamide is a functionalized amino acid. Its molecular formula is C₁₂H₁₀N₂O₂ and its molecular weight is 250.30. The chemical structure is:

Lacosamide USP is a white to light yellow powder. It is freely soluble in methanol, soluble in anhydrous ethanol, sparingly soluble in water, slightly soluble in acetonitrile practically insoluble in heptane.

11.2 Lacosamide Injection, USP $Lacosamide\ injection,\ USP\ is\ a\ clear,\ colorless,\ sterile\ solution\ containing\ 10\ mg\ lacosamide\ per\ mL\ for\ intravenous\ infusion.\ One\ 20\ mL\ vial\ contains\ 200\ mL\ vial\ conta$ mg of lacosamide drug substance. The inactive ingredients are sodium chloride (7.60 mg/mL) and water for injection. Hydrochloric acid is used for pH tment. Lacosamide injection, USP has a pH of 3.8 to 5.0.

12.1 Mechanism of Action

The precise mechanism by which lacosamide exerts its antiepileptic effects in humans remains to be fully elucidated. In vitro electrophysiological studies have shown that lacosamide selectively enhances slow inactivation of voltage-gated sodium channels, resulting in stabilization of hyperexcitable neuronal membranes and inhibition of repetitive neuronal firing.

A pharmacokinetic-pharmacodynamic (efficacy) analysis was performed based on the pooled data from the 3 efficacy trials for partial-onset seizures. Lacosamide exposure is correlated with the reduction in seizure frequency. However, doses above 400 mg/day do not appear to confer additional benefit Cardiac Electrophysiology

Electrocardiographic effects of lacosamide were determined in a double-blind, randomized clinical pharmacology trial of 247 healthy subjects. Chronic oral doses of 400 and 800 mg/day (equal to and two times the maximum daily recommended dose, respectively) were compared with placebo and a positive control (400 mg moxifloxacin). Lacosamide did not prolong QTc interval and did not have a dose-related or clinically important effect on QRS duration. Lacosamide produced a small, dose-related increase in mean PR interval. At steady-state, the time of the maximum observed mean PR interval corresponded with t... The placebo-subtracted maximum increase in PR interval (at t...) was 7.3 ms for the 400 mg/day group and 11.9 ms for the 800 mg/day group. For patients who participated in the controlled trials, the placebo-subtracted mean maximum increase in PR interval for a 400 mg/day lacosamide dose was 3.1 ms in patients with partial-onset seizures and 9.4 ms for patients with diabetic neuropathy. 12.3 Pharmacokinetics

The pharmacokinetics of lacosamide have been studied in healthy adult subjects (age range 18 to 87), adults with partial-onset seizures, adults with diabetic neuropathy, and subjects with renal and hepatic impairment. The pharmacokinetics of lacosamide are similar in healthy subjects, patients with partial-onset seizures, and patients with primary generalized tonic-clonic seizures.

Lacosamide is completely absorbed after oral administration with negligible first-pass effect with a high absolute bioavailability of approximately 100% The maximum lacosamide plasma concentrations occur approximately 1-to-4-hour post-dose after oral dosing, and elimination half-life is approximately 13 hours. Steady state plasma concentrations are achieved after 3 days of twice daily repeated administration. Pharmacokinetics of lacosamide are dose proportional (100 to 800 mg) and time invariant, with low inter- and intra-subject variability. Compared to lacosamide the major metabolite, 0desmethyl metabolite, has a longer $T_{\rm max}$ (0.5 to 12 hours) and elimination half-life (15 to 23 hours). Absorption and Bioavailability

Lacosamide is completely absorbed after oral administration. The oral bioavailability of lacosamide tablets is approximately 100%. Food does not affect the rate and extent of absorption.

After intravenous administration, C_{max} is reached at the end of infusion. The 30- and 60-minute intravenous infusions are bioequivalent to the oral tablet. For the 15-minute intravenous infusion, bioequivalence was met for AUC $_{\rm tot}$ but not for $C_{\rm ms}$. The point estimate of $C_{\rm ms}$ was 20% higher than $C_{\rm ms}$ for oral tablet and the 90% CI for $C_{\rm ms}$ exceeded the upper boundary of the bioequivalence range. $In a trial comparing the oral tablet with an oral solution containing 10\,mg/mL \, lacosamide, bioequivalence \, between both formulations \, was shown.$

A single loading dose of 200 mg approximates steady-state concentrations comparable to the 100 mg twice daily oral administration. <u>Dist</u>ribution

The volume of distribution is approximately 0.6 L/kg and thus close to the volume of total body water. Lacosamide is less than 15% bound to plasma

Lacosamide is primarily eliminated from the systemic circulation by renal excretion and biotransformation. $After intravenous administration \ of 100\,mg \ [14C] - lacosamide \ approximately \ 95\% \ of radioactivity \ administered \ was recovered in the urine \ and \ less \ than \ approximately \ 95\% \ of radioactivity \ administered \ was recovered in the urine \ and \ less \ than \ approximately \ 95\% \ of radioactivity \ administered \ was recovered in the urine \ and \ less \ than \ approximately \ 95\% \ of radioactivity \ administered \ was recovered in the urine \ and \ less \ than \ approximately \ 95\% \ of \ radioactivity \ administered \ was recovered \ in the urine \ and \ less \ than \ approximately \ 95\% \ of \ radioactivity \ administered \ was recovered \ in the urine \ and \ less \ than \ approximately \ 95\% \ of \ radioactivity \ administered \ was recovered \ in the \ urine \ and \ less \ than \ approximately \ 95\% \ of \ radioactivity \ administered \ was recovered \ in the \ urine \ and \ less \ than \ approximately \ 95\% \ of \ radioactivity \ administered \ was \ recovered \ approximately \ 95\% \ of \ radioactivity \ administered \ was \ recovered \ approximately \ 95\% \ of \ radioactivity \ administered \ was \ recovered \ approximately \ 95\% \ of \ radioactivity \ administered \ was \ recovered \ approximately \ 95\% \ of \ radioactivity \ administered \ was \ recovered \ approximately \ 95\% \ of \ radioactivity \ administered \ was \ recovered \ approximately \ 95\% \ of \ radioactivity \ administered \ was \ recovered \ approximately \ 95\% \ of \ radioactivity \ administered \ was \ recovered \ approximately \ 95\% \ of \ radioactivity \ administered \ was \ recovered \ approximately \$

0.5% in the feces. The major compounds excreted were unchanged lacosamide (approximately 40% of the dose), its 0-desmethyl metabolite (approximately 30%), and a structurally unknown polar fraction (~20%). The plasma exposure of the major human metabolite, 0-desmethyl-lacosamide, is approximately 10% of that of lacosamide. This metabolite has no known pharmacological activity.

The CYP isoforms mainly responsible for the formation of the major metabolite (O-desmethyl) are CYP3A4, CYP2C9, and CYP2C19. The elimination halflife of the unchanged drug is approximately 13 hours and is not altered by different doses, multiple dosing or intravenous administration There is no enantiomeric interconversion of lacosamide.

Specific Populations

Lacosamide and its major metabolite are eliminated from the systemic circulation primarily by renal excretion

The AUC of lacosamide was increased approximately 25% in mildly (CL_{ca} 50 to 80 mL/min) and moderately (CL_{ca} 30 to 50 mL/min) and 60% in severely (CL_{CR} \leq 30 mL/min) renally impaired patients compared to subjects with normal renal function (CL_{CR} > 80 mL/min), whereas C_{max} was unaffected. Lacosamide is effectively removed from plasma by hemodialysis. Following a 4-hour hemodialysis treatment, AUC of lacosamide is reduced by approximately 50% [see Dosage and Administration (2.4)].

Lacosamide undergoes metabolism. Subjects with moderate hepatic impairment (Child-Pugh B) showed higher plasma concentrations of

lacosamide (approximately 50 to 60% higher AUC compared to healthy subjects). The pharmacokinetics of lacosamide have not been evaluated in epatic impairment [see Dosage and Administration (2.5)]. Pediatric Patients Pediatric use information is approved for UCB, Inc.'s VIMPAT®(lacosamide) injection. However, due to UCB, Inc.'s marketing exclusivity rights,

this drug product is not labeled with that pediatric information Geriatric Patients

In the elderly (> 65 years), dose and body-weight normalized AUC and $C_{\mbox{\tiny max}}$ is about 20% increased compared to young subjects (18 to 64 years).

... Lacosamide clinical trials indicate that gender does not have a clinically relevant influence on the pharmacokinetics of lacosamide.

There are no clinically relevant differences in the pharmacokinetics of Jacosamide between Asian, Black, and Caucasian subjects.

There are no clinically relevant differences in the pharmacokinetics of lacosamide between CYP2C19 poor metabolizers and extensive netabolizers. Results from a trial in poor metabolizers (PM) (N – 4) and extensive metabolizers (EM) (N – 8) of cytochrome P450 (CYP) 2C19 showed that lacosamide plasma concentrations were similar in PMs and EMs, but plasma concentrations and the amount excreted into urine of

the O-desmethyl metabolite were about 70% reduced in PMs compared to EMs. Drug Interactions In Vitro Assessment of Drug Interactions

In vitro metabolism studies indicate that lacosamide does not induce the enzyme activity of drug metabolizing cytochrome P450 isoforms CYP1A2, 2B6, 2C9, 2C19 and 3A4. Lacosamide did not inhibit CYP1A1, 1A2, 2A6, 2B6, 2C8, 2C9, 2D6, 2E1, 3A4/5 at plasma concentrations In vitro data suggest that lacosamide has the potential to inhibit CYP2C19 at therapeutic concentrations. However, an in vivo study with

Lacosamide was not a substrate or inhibitor for P-glycoprotein Lacosamide is a substrate of CYP3A4, CYP2C9, and CYP2C19. Patients with renal or hepatic impairment who are taking strong inhibitors of

CYP3A4 and CYP2C9 may have increased exposure to Jacosamid $Since < 15\% \ of lacosamide \ is \ bound \ to \ plasma \ proteins, \ a \ clinically \ relevant \ interaction \ with \ other \ drugs \ through \ competition \ for \ protein \ binding$ sites is unlikely.

In Vivo Assessment of Drug Interactions

o <u>Effect of lacosamide on concomitant AEDs</u> Lacosamide 400 mg/day had no influence on the pharmacokinetics of 600 mg/day valproic acid and 400 mg/day carbamazepine in healthy

The placebo-controlled clinical studies in patients with partial-onset seizures showed that steady-state plasma concentrations of The process controlled clinical sculines in patients with partial controlled eaziers anowed that steady-state phasinic concentrations devertined and representations of the process of the Effect of concomitant AEDs on lacosamide

Drug-drug interaction studies in healthy subjects showed that 600 mg/day valproic acid had no influence on the pharmacokinetics of 400 mg/day lacosamide. Likewise, 400 mg/day carbamazepine had no influence on the pharmacokinetics of lacosamide in a healthy subject study. Population pharmacokinetics results in patients with partial-onset seizures showed small reductions (15% to 20% lower) in lacosamide plasma concentrations when lacosamide was coadministered with carbamazepine, phenobarbital or phenytoin.

Drug-drug interaction studies with other drugs

Digoxin There was no effect of lacosamide (400 mg/day) on the pharmacokinetics of digoxin (0.5 mg once daily) in a study in healthy subjects

There were no clinically relevant changes in met formin levels following coadministration of lacosamide (400 mg/day). Met formin (500 mg three times a day) had no effect on the pharmacokinetics of lacosamide (400 mg/day).

<u>Omeprazole</u> Omeprazole is a CYP2C19 substrate and inhibitor.

There was no effect of lacosamide (600 mg/day) on the pharmacokinetics of omeprazole (40 mg single dose) in healthy subjects. The data indicated that lacosamide had little in vivo inhibitory or inducing effect on CYP2C19. Omeprazole at a dose of 40 mg once daily had no effect on the pharmacokinetics of lacosamide (300 mg single dose). However, plasma levels of the O-desmethyl metabolite were reduced about 60% in the presence of omeprazole.

o Midazolam Midazolami si a 3A4 substrate.
There was no effect of lacosamide (200 mg single dose or repeat doses of 400 mg/day given as 200 mg BID) on the pharmacokinetics of

midazolam (single dose, 7.5 mg), indicating no inhibitory or inducing effects on CYP3A4. Oral Contraceptives

There was no influence of lacosamide (400 mg/day) on the pharmacodynamics and pharmacokinetics of an oral contraceptive containing 0.03 mg ethinylestradiol and 0.15 mg levonorgestrel in healthy subjects, except that a 20% increase in ethinylestradiol C... was observed. Warfarin

Co-administration of lacosamide (400 mg/day) with warfarin (25 mg single dose) did not result in a clinically relevant change in the pharmacokinetic and pharmacodynamic effects of warfarin in a study in healthy male subjects.

13 NONCLINICAL TOXICOLOGY 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

There was no evidence of drug related carcinogenicity in mice or rats. Mice and rats received lacosamide once daily by oral administration for 104 weeks at doses producing plasma exposures (AUC) up to approximately 1 and 3 times, respectively, the plasma AUC in humans at the maximun human dose (MRHD) of 400 mg/day.

Lacosamide was negative in an in vitro Ames test and an in vivo mouse micronucleus assay. Lacosamide induced a positive response in the in vitro mouse <u>Fertility.</u>

No adverse effects on male or female fertility or reproduction were observed in rats at doses producing plasma exposures (AUC) up to approximately

14 CLINICAL STUDIES

14.1 Monotherapy in Patients with Partial-Onset Seizures

The efficacy of lacosamide in monotherapy was established in a historical-control, multicenter, randomized trial that included 425 patients, age 16 to 70 years, with partial-onset seizures (Study 1). To be included in Study 1, patients were required to be taking stable doses of 1 or 2 marketed antispileptic drugs. This treatment continued into the 8 week baseline period. To remain in the study, patients were required to have at least 2 partial-onset seizures per 28 days during the 8 week baseline period. The baseline period was followed by a 3 week titration period, during which lacosamide was added to the ongoing antiepileptic regimen. This was followed by a 16-week maintenance period (i.e., a 6-week withdrawal period for background antispileptic drugs, followed by a 10-week monotherapy period). Patients were randomized 3 to 1 to receive lacosamide 400 mg/day or lacosamide 300 mg/day. Treatment assignments were blinded. Response to treatment was based upon a comparison of the number of patients who met exit criteria

lacosamide remained below the lower 95% prediction limit of 65% derived from the historical control data. The exit criteria were one or more of the following: (1) doubling of average monthly seizure frequency during any 28 consecutive days. (2) doubling of highest consecutive 2-day seizure frequency, (3) occurrence of a single generalized tonic clonic seizure, (4) clinically significant prolongation of worsening of overall seizure duration, frequency, type or pattern considered by the investigator to require trial discontinuation, (5) status epilepticus or new onset of serial/cluster seizures. The study population profile appeared comparable to that of the historical control population For the lacosamide 400 mg/day group, the estimate of the percentage of patients meeting at least 1 exit criterion was 30% (95% CI: 25%, 36%). The

during the maintenance phase, compared to historical controls. The historical control consisted of a pooled analysis of the control groups from 8 studies of similar design, which utilized a sub-therapeutic dose of an antiepileptic drug. Statistical superiority to the historical control was considered to be demonstrated if the upper limit from a 2-sided 95% confidence interval for the percentage of patients meeting exit criteria in patients receiving

upper limit of the 2-sided 95% CI (36%) was below the threshold of 65% derived from the historical control data, meeting the pre-specified criteria fo efficacy. Lacosamide 300 mg/day also met the pre-specified criteria for efficacy. 14.2 Adjunctive Therapy in Patients with Partial-Onset Seizures The efficacy of lacosamide as adjunctive therapy in partial-onset seizures was established in three 12-week, randomized, double-blind, placebocontrolled, multicenter trials in adult patients (Study 2, Study 3, and Study 4). Enrolled patients had partial-onset seizures with or without secondary generalization, and were not adequately controlled with 1 to 3 concomitant AEDs. During an 8-week baseline period, patients were required to have an average of 2-4 partial-onset seizures per 28 days with no seizure-free period exceeding 21 days. In these 3 trials, patients had a mean duration of epilepsy of 24 years and a median baseline seizure frequency ranging from 10 to 17 per 28 days. 84% of patients were taking 2 to 3 concomitant AEDs

with or without concurrent vagal nerve stimulation. Study 2 compared doses of lacosamide 200, 400, and 600 mg/day with placebo, Study 3 compared doses of lacosamide 400 and 600 mg/day with placebo. Study 4 compared doses of lacosamide 200 and 400 mg/day with placebo. In all three trials, following an 8-week baseline phase to establish baseline seizure frequency prior to randomization, patients were randomized and titrated to the randomized dose (a 1-step back-titration of lacosamide 100 molday or placebo was allowed in the case of intolerable adverse reactions at the end of the titration phase. During the titration phase, in all 3

adjunctive therapy trials, treatment was initiated at 100 mg/day (50 mg twice daily) and increased in weekly increments of 100 mg/day to the target dose. The titration phase lasted 6 weeks in Study 2 and Study 3, and 4 weeks in Study 4. In all three trials, the titration phase was followed by a nce phase that lasted 12 weeks, during which patients were to remain on a stable dose of lacosamide A reduction in 28-day seizure frequency (baseline to maintenance phase), as compared to the placeho group, was the primary variable in all three adjunctive therapy trials. A statistically significant effect was observed with lacosamide treatment (Figure 1) at doses of 200 mg/day (Study 4),

400 mg/day (Studies 2, 3, and 4), and 600 mg/day (Studies 2 and 3). tions of lacosamide demonstrate no important differences in seizure control as a function of gender or race, although data on race was

limited (about 10% of patients were non-Caucasian)

Figure 1- Median Percent Reduction in Seizure Frequency per 28 days from Baseline to the Maintenance Phase by Dose





Figure 2 presents the percentage of patients (X-axis) with a percent reduction in partial seizure frequency (responder rate) from baseline to the night of presents up becoming or placetar (x-axis) which a percent reduction in placeta sector requester (response to the percent of the presented on the Y-axis and the percent of the percent of the Y-axis indicates an improvement from baseline (i.e., a decrease in seizure frequency), while a negative value indicates a worsening from baseline (i.e., an increase in seizure frequency). Thus, in a display of this type, a curve for an effective treatment is shifted to the left of the curve for placeho. The proportion of nationals achieving any particular level of reduction in seizure frequency was consistently higher for the lacosamide groups, compared to the placebo group. For example, 40% of patients randomized to lacosamide (400 mg/day) experienced a 50% or greater reduction in seizure frequency, compared to 23% of patients randomized to placebo. Patients with an increase in seizure frequency > 100% are represented on the Y-axis as equal to or greater than -100%



Pediatric use information is approved for UCB, Inc.'s VIMPAT® (lacosamide) injection. However, due to UCB, Inc.'s marketing exclusivity rights, this

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied Lacosamide Injection LISP

 $200\,mg/20\,mL\,is\,a\,clear, colorless\,sterile\,solution\,supplied\,in\,20\,mL\,colorless\,single\cdot dose\,glass\,vials.$ 200 mg/20 mL vial in cartons of 10 vials NDC 31722-203-31 16.2 Storage and Handling

Store at 20 °C to 25 °C (68 °F to 77 °F); excursions permitted between 15 °C to 30 °C (59 °F to 86 °F). [See USP Controlled Room Temperature]

17 PATIENT COUNSELING INFORMATION Advise the patient or caregiver to read the FDA-approved patient labeling (Medication Guide). The Medication Guide accompanies the product and can also be accessed by calling 1-866-495-1995.

Suicidal Thinking and Behavior Patients, their caregivers, and families should be counseled that AEDs, including lacosamide, may increase the risk of suicidal thoughts and behavior and should be advised of the need to be alert for the emergence or worsening of symptoms of depression, any unusual changes in mood or behavior, or the emergence of suicidal thoughts, behavior, or thoughts about self-harm. Behaviors of concern should be reported immediately to healthcare providers /see

Warnings and Precautions (5.1)]. Dizziness and Ataxia Patients should be counseled that lacosamide use may cause dizziness, double vision, abnormal coordination and balance, and somnolence. Patients Facients should be considered that recosame use may be used success, outlier vision, another account and unanner, and shindle hardware account taking lacosamide should be advised not to drive, operate complex machinery, or engage in other hazardous activities until they have become accust to any such effects associated with lacosamide (see Warnings and Precautions (5.2)).

Cardiac Rhythm and Conduction Abnormalities Patients should be counseled that lacosamide is associated with electrocardiographic changes that may predispose to irregular heart beat and syncope. Cardiac arrest has been reported. This risk is increased in patients with underlying cardiovascular disease, with heart conduction problems, or who are taking other medications that affect the heart. Patients should be made aware of and report cardiac signs or symptoms to their healthcare provider right away. Patients who develop syncope should lay down with raised legs and contact their health care provider [see Warnings and Precautions (5.3]]

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Multi-Organ Hypersensitivity Patients should be aware that lacosamide may cause serious hypersensitivity reactions affecting multiple organs such as the liver and kidney. Lacosamide should be discontinued if a serious hypersensitivity reaction is suspected. Patients should also be instructed to report promptly to their physicians any symptoms of liver toxicity (e.g., fatigue, jaundice, dark urine) [see Warnings and Precautions (5.6)].

Pregnancy Registry Advise patients to notify their healthcare provider if they become pregnant or intend to become pregnant during lacosamide therapy. Encourage patients to enroll in the North American Antiepileptic Drug (NAAED) pregnancy registry if they become pregnant. This registry is collecting information about the

safety of AEDs during pregnancy /see Use in Specific Populations (8.1)/. Advise breastfeeding women using lacosamide to monitor infants for excess sleepiness and to seek medical care if they notice this sign /see Use in

Specific Populations (8.2)].

Camber Pharmaceuticals, Inc. Piscataway, NJ 08854

Manufactured for:

∧ Aspiro Aspiro Pharma Limited Survey No. 321, Biotech Park, Phase – III Karkapatla Village, Markook (Mandal) Siddipet, Telangana-502281, India.

21002220