

60 mm

**Dosage:** See package insert.

Net contents after constitution: 100 mL, containing 5 grams valganciclovir free base.

Deliverable volume after constitution: 88 mL.

**To the pharmacist:** Prepare the solution at the time of dispensing by measuring 91 mL of purified water. Add approximately half the total amount of water and shake closed bottle well for about 1 minute.

Add remainder of water and shake closed bottle well for about 1 minute. Open and push bottle adapter into the neck of the bottle and close with child resistant cap.

The constituted oral solution should be used within 49 days of preparation, write the Discard Date of the constituted oral solution on the bottle label. Dispense with patient instructions and 2 oral dispensers.

**Store dry powder at 20°C to 25°C (68°F to 77°F); excursions are permitted to 15°C to 30°C (59°F to 86°F) [see USP controlled room temperature].**

**Store constituted solution under refrigeration at 2°C to 8°C (36°F to 46°F). Do not freeze.**

Store in the original bottle. Keep the bottle tightly closed.

Avoid direct contact with powder and constituted solution.

Discard any unused portion 49 days after constitution.



NDC 31722-837-10

## Valganciclovir for Oral Solution

50 mg/mL

Each mL of constituted oral solution contains 50 mg valganciclovir free base.

Rx only

100 mL (3.4 fl oz)

Un varnish Area for  
Batch Coding  
40x25 mm

2102149

Constitute Date: .....

Discard Date: .....

U.S. Contact Number: 1-866-495-1995

Mfg. Lic. No.: 24/MD/TS/2016/FF/G

Manufactured by:  
Camber Pharmaceuticals, Inc.  
Piscataway, NJ 08854.

By: Annora Pharma Pvt. Ltd.  
Sangareddy - 502313, Telangana, India.



Size : 120 x 60 mm

Colours : **PANTONE 1805 C** **PANTONE 2736 C** **PANTONE Black C**

Non Printing colours:   Die cut

120 mm