

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use LISDEXAMFETAMINE DIMESYLATE CAPSULES safely and effectively. See full prescribing information for LISDEXAMFETAMINE DIMESYLATE CAPSULES.

### LISDEXAMFETAMINE DIMESYLATE capsules, for oral use, CII

Initial U.S. Approval: 2007

| WARNING: ABUSE, MISUSE, AND ADDICTION                                                                                                                                                                                                                                                                                                                                                                                                                    |                                                                     |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------|
|                                                                                                                                                                                                                                                                                                                                                                                                                                                          | <i>See full prescribing information for complete boxed warning.</i> |
| Lisdexamfetamine dimesylate capsules has a high potential for abuse and misuse, which can lead to the development of a substance use disorder, including addiction. Misuse and abuse of CNS stimulants, including lisdexamfetamine dimesylate capsules, can result in overdose and death (5.1, 9.2, 10).                                                                                                                                                 |                                                                     |
| <ul style="list-style-type: none"><li>• Before prescribing lisdexamfetamine dimesylate capsules, assess each patient's risk for abuse, misuse, and addiction.</li><li>• Educate patients and their families about these risks, proper storage of the drug, and proper disposal of any unused drug.</li><li>• Throughout treatment, reassess each patient's risk and frequently monitor for signs and symptoms of abuse, misuse, and addiction.</li></ul> |                                                                     |

### RECENT MAJOR CHANGES

|                                |         |
|--------------------------------|---------|
| Indications and Usage (1)      | 09/2025 |
| Warnings and Precautions (5.5) | 09/2025 |

Lisdexamfetamine dimesylate capsules are a central nervous system (CNS) stimulant indicated for the treatment of (1):

- Attention Deficit Hyperactivity Disorder (ADHD) in adults and pediatric patients 6 years and older
- Moderate to severe binge eating disorder (BED) in adults

**Limitations of Use:**

- The use of lisdexamfetamine dimesylate capsules are not recommended in pediatric patients younger than 6 years of age because they have higher plasma exposure and a higher incidence of adverse reactions (e.g., weight loss) than patients 6 years and older at the same dosage (5.5, 8, 4).
- Lisdexamfetamine dimesylate capsules are not indicated for weight loss. Use of other sympathomimetic drugs for weight loss has been associated with serious cardiovascular adverse events. The safety and effectiveness of lisdexamfetamine dimesylate capsules for the treatment of obesity have not been established (5.2).

| DOSAGE AND ADMINISTRATION                                    |                     |                       |                        |               |
|--------------------------------------------------------------|---------------------|-----------------------|------------------------|---------------|
| Indicated Population                                         | Initial Dose        | Titration Schedule    | Recommended Dose       | Maximum Dose  |
| ADHD (Adults and pediatric patients 6 years and older) (2,2) | 30 mg every morning | 10 mg or 20 mg weekly | 30 mg to 70 mg per day | 70 mg per day |
| BED (Adults) (2,3)                                           | 30 mg every morning | 20 mg weekly          | 50 mg to 70 mg per day | 70 mg per day |

- Prior to treatment, assess for presence of cardiac disease (2,4)
- Severe renal impairment: Maximum dose is 50 mg daily (2,5)

### FULL PRESCRIBING INFORMATION: CONTENTS\*

#### WARNING: ABUSE, MISUSE, AND ADDICTION

#### 1 INDICATIONS AND USAGE

#### 2 DOSAGE AND ADMINISTRATION

- 2.1 Pretreatment Screening
- 2.2 General Administration Information
- 2.3 Dosage for Treatment of ADHD
- 2.4 Dosage for Treatment of Moderate to Severe BED in Adults
- 2.5 Dosage in Patients with Renal Impairment
- 2.6 Dosage Modifications due to Drug Interactions

#### 3 DOSAGE FORMS AND STRENGTHS

#### 4 CONTRAINDICATIONS

#### 5 WARNINGS AND PRECAUTIONS

- 5.1 Abuse, Misuse, and Addiction
- 5.2 Risks to Patients with Serious Cardiac Disease
- 5.3 Increased Blood Pressure and Heart Rate
- 5.4 Psychiatric Adverse Reactions
- 5.5 Long-Term Suppression of Growth in Pediatric Patients
- 5.6 Peripheral Vascuopathy, including Raynaud's Phenomenon
- 5.7 Serotonin Syndrome
- 5.8 Motor and Verbal Tics, and Worsening of Tourette's Syndrome

#### 6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

#### 7 DRUG INTERACTIONS

- 7.1 Drugs Having Clinically Important Interactions with Amphetamines
- 7.2 Drugs Having No Clinically Important Interactions with Lisdexamfetamine Dimesylate Capsules

#### 8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy

#### 9 OVERDOSAGE

#### 10 CLINICAL PHARMACOLOGY

#### 11 DESCRIPTION

#### 12 NONCLINICAL TOXICOLOGY

#### 13 CARCINOGENESIS, MUTAGENESIS, AND IMPAIRMENT OF FERTILITY

#### 14 CLINICAL STUDIES

#### 15 HOW SUPPLIED/STORAGE AND HANDLING

#### 16 PATIENT COUNSELING INFORMATION

#### 17 REFERENCES

#### 18 SUPPLEMENTAL INFORMATION

#### 19 PATENT INFORMATION

#### 20 TRADEMARKS

#### 21 OTHER INFORMATION

#### 22 SUPPLEMENTAL INFORMATION

#### 23 SUPPLEMENTAL INFORMATION

#### 24 SUPPLEMENTAL INFORMATION

#### 25 SUPPLEMENTAL INFORMATION

#### 26 SUPPLEMENTAL INFORMATION

#### 27 SUPPLEMENTAL INFORMATION

#### 28 SUPPLEMENTAL INFORMATION

#### 29 SUPPLEMENTAL INFORMATION

#### 30 SUPPLEMENTAL INFORMATION

#### 31 SUPPLEMENTAL INFORMATION

#### 32 SUPPLEMENTAL INFORMATION

#### 33 SUPPLEMENTAL INFORMATION

#### 34 SUPPLEMENTAL INFORMATION

#### 35 SUPPLEMENTAL INFORMATION

#### 36 SUPPLEMENTAL INFORMATION

#### 37 SUPPLEMENTAL INFORMATION

#### 38 SUPPLEMENTAL INFORMATION

#### 39 SUPPLEMENTAL INFORMATION

#### 40 SUPPLEMENTAL INFORMATION

#### 41 SUPPLEMENTAL INFORMATION

#### 42 SUPPLEMENTAL INFORMATION

#### 43 SUPPLEMENTAL INFORMATION

#### 44 SUPPLEMENTAL INFORMATION

#### 45 SUPPLEMENTAL INFORMATION

#### 46 SUPPLEMENTAL INFORMATION

#### 47 SUPPLEMENTAL INFORMATION

#### 48 SUPPLEMENTAL INFORMATION

#### 49 SUPPLEMENTAL INFORMATION

#### 50 SUPPLEMENTAL INFORMATION

#### 51 SUPPLEMENTAL INFORMATION

#### 52 SUPPLEMENTAL INFORMATION

#### 53 SUPPLEMENTAL INFORMATION

#### 54 SUPPLEMENTAL INFORMATION

#### 55 SUPPLEMENTAL INFORMATION

#### 56 SUPPLEMENTAL INFORMATION

#### 57 SUPPLEMENTAL INFORMATION

#### 58 SUPPLEMENTAL INFORMATION

#### 59 SUPPLEMENTAL INFORMATION

#### 60 SUPPLEMENTAL INFORMATION

#### 61 SUPPLEMENTAL INFORMATION

#### 62 SUPPLEMENTAL INFORMATION

#### 63 SUPPLEMENTAL INFORMATION

#### 64 SUPPLEMENTAL INFORMATION

#### 65 SUPPLEMENTAL INFORMATION

#### 66 SUPPLEMENTAL INFORMATION

#### 67 SUPPLEMENTAL INFORMATION

#### 68 SUPPLEMENTAL INFORMATION

#### 69 SUPPLEMENTAL INFORMATION

#### 70 SUPPLEMENTAL INFORMATION

#### 71 SUPPLEMENTAL INFORMATION

#### 72 SUPPLEMENTAL INFORMATION

#### 73 SUPPLEMENTAL INFORMATION

#### 74 SUPPLEMENTAL INFORMATION

#### 75 SUPPLEMENTAL INFORMATION

#### 76 SUPPLEMENTAL INFORMATION

#### 77 SUPPLEMENTAL INFORMATION

#### 78 SUPPLEMENTAL INFORMATION

#### 79 SUPPLEMENTAL INFORMATION

#### 80 SUPPLEMENTAL INFORMATION

#### 81 SUPPLEMENTAL INFORMATION

#### 82 SUPPLEMENTAL INFORMATION

#### 83 SUPPLEMENTAL INFORMATION

#### 84 SUPPLEMENTAL INFORMATION

#### 85 SUPPLEMENTAL INFORMATION

#### 86 SUPPLEMENTAL INFORMATION

#### 87 SUPPLEMENTAL INFORMATION

#### 88 SUPPLEMENTAL INFORMATION

#### 89 SUPPLEMENTAL INFORMATION

#### 90 SUPPLEMENTAL INFORMATION

#### 91 SUPPLEMENTAL INFORMATION

#### 92 SUPPLEMENTAL INFORMATION

#### 93 SUPPLEMENTAL INFORMATION

#### 94 SUPPLEMENTAL INFORMATION

#### 95 SUPPLEMENTAL INFORMATION

#### 96 SUPPLEMENTAL INFORMATION

#### 97 SUPPLEMENTAL INFORMATION

#### 98 SUPPLEMENTAL INFORMATION

#### 99 SUPPLEMENTAL INFORMATION

#### 100 SUPPLEMENTAL INFORMATION

#### 101 SUPPLEMENTAL INFORMATION

#### 102 SUPPLEMENTAL INFORMATION

#### 103 SUPPLEMENTAL INFORMATION

#### 104 SUPPLEMENTAL INFORMATION

#### 105 SUPPLEMENTAL INFORMATION

#### 106 SUPPLEMENTAL INFORMATION

#### 107 SUPPLEMENTAL INFORMATION

#### 108 SUPPLEMENTAL INFORMATION

#### 109 SUPPLEMENTAL INFORMATION

#### 110 SUPPLEMENTAL INFORMATION

#### 111 SUPPLEMENTAL INFORMATION

#### 112 SUPPLEMENTAL INFORMATION

#### 113 SUPPLEMENTAL INFORMATION

#### 114 SUPPLEMENTAL INFORMATION

#### 115 SUPPLEMENTAL INFORMATION

#### 116 SUPPLEMENTAL INFORMATION

#### 117 SUPPLEMENTAL INFORMATION

#### 118 SUPPLEMENTAL INFORMATION

#### 119 SUPPLEMENTAL INFORMATION

#### 120 SUPPLEMENTAL INFORMATION

#### 121 SUPPLEMENTAL INFORMATION

#### 122 SUPPLEMENTAL INFORMATION

#### 123 SUPPLEMENTAL INFORMATION

#### 124 SUPPLEMENTAL INFORMATION

#### 125 SUPPLEMENTAL INFORMATION

#### 126 SUPPLEMENTAL INFORMATION

#### 127 SUPPLEMENTAL INFORMATION

#### 128 SUPPLEMENTAL INFORMATION

#### 129 SUPPLEMENTAL INFORMATION

#### 130 SUPPLEMENTAL INFORMATION

#### 131 SUPPLEMENTAL INFORMATION

#### 132 SUPPLEMENTAL INFORMATION

#### 133 SUPPLEMENTAL INFORMATION

#### 134 SUPPLEMENTAL INFORMATION

#### 135 SUPPLEMENTAL INFORMATION

#### 136 SUPPLEMENTAL INFORMATION

#### 137 SUPPLEMENTAL INFORMATION

#### 138 SUPPLEMENTAL INFORMATION

#### 139 SUPPLEMENTAL INFORMATION

#### 140 SUPPLEMENTAL INFORMATION

#### 141 SUPPLEMENTAL INFORMATION

#### 142 SUPPLEMENTAL INFORMATION

#### 143 SUPPLEMENTAL INFORMATION

#### 144 SUPPLEMENTAL INFORMATION

#### 145 SUPPLEMENTAL INFORMATION

#### 146 SUPPLEMENTAL INFORMATION

#### 147 SUPPLEMENTAL INFORMATION

#### 148 SUPPLEMENTAL INFORMATION

#### 149 SUPPLEMENTAL INFORMATION

#### 150 SUPPLEMENTAL INFORMATION

#### 151 SUPPLEMENTAL INFORMATION

#### 152 SUPPLEMENTAL INFORMATION

#### 153 SUPPLEMENTAL INFORMATION

#### 154 SUPPLEMENTAL INFORMATION

#### 155 SUPPLEMENTAL INFORMATION

#### 156 SUPPLEMENTAL INFORMATION

#### 157 SUPPLEMENTAL INFORMATION

#### 158 SUPPLEMENTAL INFORMATION

#### 159 SUPPLEMENTAL INFORMATION

#### 160 SUPPLEMENTAL INFORMATION

#### 161 SUPPLEMENTAL INFORMATION

#### 162 SUPPLEMENTAL INFORMATION

#### 163 SUPPLEMENTAL INFORMATION

#### 164 SUPPLEMENTAL INFORMATION

#### 165 SUPPLEMENTAL INFORMATION

#### 166 SUPPLEMENTAL INFORMATION

#### 167 SUPPLEMENTAL INFORMATION

#### 168 SUPPLEMENTAL INFORMATION

#### 169 SUPPLEMENTAL INFORMATION

#### 170 SUPPLEMENTAL INFORMATION

#### 171 SUPPLEMENTAL INFORMATION

#### 172 SUPPLEMENTAL INFORMATION

#### 173 SUPPLEMENTAL INFORMATION

#### 174 SUPPLEMENTAL INFORMATION

#### 175 SUPPLEMENTAL INFORMATION

#### 176 SUPPLEMENTAL INFORMATION

#### 177 SUPPLEMENTAL INFORMATION

#### 178 SUPPLEMENTAL INFORMATION

#### 179 SUPPLEMENTAL INFORMATION

#### 180 SUPPLEMENTAL INFORMATION

#### 181 SUPPLEMENTAL INFORMATION

#### 182 SUPPLEMENTAL INFORMATION

#### 183 SUPPLEMENTAL INFORMATION

#### 184 SUPPLEMENTAL INFORMATION

#### 185 SUPPLEMENTAL INFORMATION

#### 186 SUPPLEMENTAL INFORMATION

#### 187 SUPPLEMENTAL INFORMATION

#### 188 SUPPLEMENTAL INFORMATION

#### 189 SUPPLEMENTAL INFORMATION

#### 190 SUPPLEMENTAL INFORMATION

#### 191 SUPPLEMENTAL INFORMATION

#### 192 SUPPLEMENTAL INFORMATION

#### 193 SUPPLEMENTAL INFORMATION

#### 194 SUPPLEMENTAL INFORMATION

#### 195 SUPPLEMENTAL INFORMATION

#### 196 SUPPLEMENTAL INFORMATION

#### 197 SUPPLEMENTAL INFORMATION

#### 198 SUPPLEMENTAL INFORMATION

#### 199 SUPPLEMENTAL INFORMATION

#### 200 SUPPLEMENTAL INFORMATION

#### 201 SUPPLEMENTAL INFORMATION

#### 202 SUPPLEMENTAL INFORMATION

#### 203 SUPPLEMENTAL INFORMATION

#### 204 SUPPLEMENTAL INFORMATION

#### 205 SUPPLEMENTAL INFORMATION

#### 206 SUPPLEMENTAL INFORMATION

#### 207 SUPPLEMENTAL INFORMATION

#### 208 SUPPLEMENTAL INFORMATION

#### 209 SUPPLEMENTAL INFORMATION

#### 210 SUPPLEMENTAL INFORMATION

#### 211 SUPPLEMENTAL INFORMATION

#### 212 SUPPLEMENTAL INFORMATION

#### 213 SUPPLEMENTAL INFORMATION

#### 214 SUPPLEMENTAL INFORMATION

#### 215 SUPPLEMENTAL INFORMATION

#### 216 SUPPLEMENTAL INFORMATION

#### 217 SUPPLEMENTAL INFORMATION

#### 218 SUPPLEMENTAL INFORMATION

#### 219 SUPPLEMENTAL INFORMATION

#### 220 SUPPLEMENTAL INFORMATION

#### 221 SUPPLEMENTAL INFORMATION

#### 222 SUPPLEMENTAL INFORMATION

#### 223 SUPPLEMENTAL INFORMATION

#### 224 SUPPLEMENTAL INFORMATION

#### 225 SUPPLEMENTAL INFORMATION

#### 226 SUPPLEMENTAL INFORMATION

#### 227 SUPPLEMENTAL INFORMATION

#### 228 SUPPLEMENTAL INFORMATION

#### 229 SUPPLEMENTAL INFORMATION

#### 230 SUPPLEMENTAL INFORMATION

#### 231 SUPPLEMENTAL INFORMATION

#### 232 SUPPLEMENTAL INFORMATION

#### 233 SUPPLEMENTAL INFORMATION

#### 234 SUPPLEMENTAL INFORMATION

#### 235 SUPPLEMENTAL INFORMATION

#### 236 SUPPLEMENTAL INFORMATION

#### 237 SUPPLEMENTAL INFORMATION

#### 238 SUPPLEMENTAL INFORMATION

#### 239 SUPPLEMENTAL INFORMATION

#### 240 SUPPLEMENTAL INFORMATION

#### 241 SUPPLEMENTAL INFORMATION

#### 242 SUPPLEMENTAL INFORMATION

#### 243 SUPPLEMENTAL INFORMATION

#### 244 SUPPLEMENTAL INFORMATION



Width: 17.0"  
Length: 22.0"  
Fold: 1.375" x 1.375"

HIGHLIGHTS OF PRESCRIBING INFORMATION  
These highlights do not include all the information needed to use LIDEXAMFETAMINE DIMESYLATE CAPSULES safely and effectively. See full prescribing information for LIDEXAMFETAMINE DIMESYLATE CAPSULES.

LIDEXAMFETAMINE DIMESYLATE capsules, for oral use, CII  
Initial U.S. Approval: 2007

**WARNING: ABUSE, MISUSE, AND ADDICTION**  
See full prescribing information for complete boxed warning.  
Lisdexamfetamine dimesylate capsules has a high potential for abuse and misuse, which can lead to the development of a substance use disorder, including addiction. Misuse and abuse of CNS stimulants, including lisdexamfetamine dimesylate capsules, can result in overdose and death (5.1, 8.2, 10).  
• Before prescribing lisdexamfetamine dimesylate capsules, assess each patient's risk for abuse, misuse, and addiction.  
• Educate patients and their families about these risks, proper storage of the drug, and proper disposal of any unused drug.  
• Throughout treatment, reassess each patient's risk and frequently monitor for signs and symptoms of abuse, misuse, and addiction.

RECENT MAJOR CHANGES

Indications and Usage (1) 09/2025  
Warnings and Precautions (5.5) 09/2025

INDICATIONS AND USAGE

Lisdexamfetamine dimesylate capsules are a central nervous system (CNS) stimulant indicated for the treatment of (1):  
• Attention Deficit Hyperactivity Disorder (ADHD) in adults and pediatric patients 6 years and older  
• Moderate to severe binge eating disorder (BED) in adults

Limitations of Use:

- The use of lisdexamfetamine dimesylate capsules are not recommended in pediatric patients younger than 6 years of age because they had higher plasma exposure and a higher incidence of adverse reactions (e.g., weight loss) than patients 6 years and older at the same dosage (5.5, 8.4)
- Lisdexamfetamine dimesylate capsules are not indicated for weight loss. Use of any sympathomimetic drug for weight loss has been associated with serious cardiovascular adverse events. The safety and effectiveness of lisdexamfetamine dimesylate capsules for the treatment of obesity have not been established (5.2).

DOSEAGE AND ADMINISTRATION

| Indicated Population                                       | Initial Dose        | Titration Schedule    | Recommended Dose       | Maximum Dose  |
|------------------------------------------------------------|---------------------|-----------------------|------------------------|---------------|
| ADHD Adults and pediatric patients 6 years and older (2,2) | 30 mg every morning | 10 mg or 20 mg weekly | 30 mg to 70 mg per day | 70 mg per day |
| BED Adults (2,3)                                           | 30 mg every morning | 20 mg weekly          | 50 mg to 70 mg per day | 70 mg per day |

- Prior to treatment, assess for presence of cardiac disease (2,4)
- Severe renal impairment: Maximum dose is 50 mg/day (2,5)

FULL PRESCRIBING INFORMATION: CONTENTS

WARNING: ABUSE, MISUSE, AND ADDICTION

1 INDICATIONS AND USAGE

2 DOSEAGE AND ADMINISTRATION

- 2.1 Prefreatment Screening
- 2.2 General Administration Information
- 2.3 Dosage for Treatment of ADHD
- 2.4 Dosage for Treatment of Moderate to Severe BED in Adults
- 2.5 Dosage in Patients with Renal Impairment
- 2.6 Dosage Modifications due to Drug Interactions

3 DOSEAGE FORMS AND STRENGTHS

- 3.1 Abuse, Misuse, and Addiction
- 3.2 Risks to Patients with Serious Cardiac Disease
- 3.3 Increased Blood Pressure and Heart Rate
- 3.4 Psychiatric Adverse Reactions

- 3.5 Long-Term Suppression of Growth in Pediatric Patients
- 3.6 Peripheral Vascopathy, including Raynaud's Phenomenon
- 3.7 Serotonin Syndrome
- 3.8 Motor and Verbal Tics, and Worsening of Tourette's Syndrome

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Drugs Having Clinically Important Interactions with Amphetamines
- 7.2 Drugs Having No Clinically Important Interactions with Lisdexamfetamine Dimesylate Capsules

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy

FULL PRESCRIBING INFORMATION

WARNING: ABUSE, MISUSE, AND ADDICTION

Lisdexamfetamine dimesylate capsules has a high potential for abuse and misuse, which can lead to the development of a substance use disorder, including addiction. Misuse and abuse of CNS stimulants, including lisdexamfetamine dimesylate capsules, can result in overdose and death (5.1, 8.2, 10).  
• Before prescribing lisdexamfetamine dimesylate capsules, assess each patient's risk for abuse, misuse, and addiction. Educate patients and their families about these risks, proper storage of the drug, and proper disposal of any unused drug. Throughout lisdexamfetamine dimesylate capsules treatment, reassess each patient's risk of abuse, misuse, and addiction and frequently monitor for signs and symptoms of abuse, misuse, and addiction (See **Warnings and Precautions (5.1, 8.2, 10)**, **Drug Abuse and Dependence (8.2)**).

1 INDICATIONS AND USAGE

Lisdexamfetamine dimesylate capsules are indicated for the treatment of:  
• Attention Deficit Hyperactivity Disorder (ADHD) in adults and pediatric patients 6 years and older (See *Clinical Studies (14.1)*)  
• Moderate to severe binge eating disorder (BED) in adults (See *Clinical Studies (14.2)*)

Limitations of Use:

- The use of lisdexamfetamine dimesylate capsules are not recommended in pediatric patients younger than 6 years of age because they had higher plasma exposure and a higher incidence of adverse reactions (e.g., weight loss) than patients 6 years and older at the same dosage (5.5, 8.4)
- Lisdexamfetamine dimesylate capsules are not indicated or recommended for weight loss. Use of any sympathomimetic drug for weight loss has been associated with serious cardiovascular adverse events. The safety and effectiveness of lisdexamfetamine dimesylate capsules for the treatment of obesity have not been established (See **Warnings and Precautions (5.2)**).

2 DOSEAGE AND ADMINISTRATION

2.1 Prefreatment Screening

Prior to treating patients with lisdexamfetamine dimesylate capsules, assess:  
• For the presence of cardiac disease (e.g., perform a careful history, family history of sudden death or ventricular arrhythmia, and physical exam) (See **Warnings and Precautions (5.2)**).

• the family history and clinically evaluate patients for motor or verbal tics or Tourette's syndrome before initiating lisdexamfetamine dimesylate capsules (See **Warnings and Precautions (5.8)**).

2.2 General Administration Information

Take lisdexamfetamine dimesylate capsules only in the morning with or without food; avoid afternoon doses because of the potential for insomnia. Lisdexamfetamine dimesylate capsules may be administered in one of the following ways:

**Information for lisdexamfetamine dimesylate capsules:**  
• Swallow lisdexamfetamine dimesylate capsules whole, or  
• Open capsules, empty and mix the entire contents with yogurt, water, or orange juice. If the contents of the capsule include any compacted powder, a spoon may be used to break apart the powder. The contents should be mixed until completely dispersed. Consume the entire mixture immediately. It should not be stored. The active ingredient dissolves completely once dispersed; however, a film containing the inactive ingredients may remain in the glass or container once the mixture is consumed.

Lisdexamfetamine dimesylate capsules can be substituted with lisdexamfetamine dimesylate chewable tablets on a unit per unit or mg per mg basis (for example, 30 mg capsules for 30 mg chewable tablets) (See *Clinical Pharmacology (12.3)*).

Do not take anything less than one capsule per day. A single dose should not be divided.

2.3 Dosage for Treatment of ADHD

The recommended starting dosage in adults and pediatric patients 6 years and older is 30 mg once daily in the morning. Dosage may be adjusted in increments of 10 mg or 20 mg at approximately weekly intervals up to maximum recommended dosage of 70 mg once daily (See *Clinical Studies (14.1)*).

2.4 Dosage for Treatment of Moderate to Severe BED in Adults

The recommended starting dosage in adults is 30 mg once daily to be titrated in increments of 20 mg at approximately weekly intervals to achieve the recommended target dose of 50 mg to 70 mg once daily. The maximum recommended dose is 70 mg once daily (See *Clinical Studies (14.2)*).

2.5 Dosage in Patients with Renal Impairment

In patients with severe renal impairment (GFR 15 to <30 mL/min/1.73 m<sup>2</sup>), the maximum dosage should not exceed 50 mg once daily in patients with end stage renal disease (ESRD, GFR <15 mL/min/1.73 m<sup>2</sup>), the maximum recommended dosage is 30 mg once daily (See *See Use in Specific Populations (8.4)*).

**2.6 Dosage Modifications due to Drug Interactions**  
Agents that alter urinary pH can impact urinary excretion and alter blood levels of amphetamine. Acidifying agents (e.g., ascorbic acid) decrease blood levels of amphetamine. Alkalinizing agents (e.g., sodium bicarbonate) increase blood levels. Adjust lisdexamfetamine dimesylate capsules dosage accordingly (See *Drug Interactions (7.1)*).

3 DOSEAGE FORMS AND STRENGTHS

**Lisdexamfetamine dimesylate capsules:**  
• Capsules 10 mg: Hard Gelatin Capsule Shell Size "3" Pink Opaque Cap imprinted with AC in Black ink and Pink Opaque body imprinted with 10 in Black ink filled with White to Off-white powder.  
• Capsules 20 mg: Hard Gelatin Capsule Shell Size "3" Ivory Opaque Cap imprinted with AC in Black ink and Ivory Opaque body imprinted with 20 in Black ink filled with White to Off-white powder.  
• Capsules 30 mg: Hard Gelatin Capsule Shell Size "3" Orange Opaque Cap imprinted with AC in Black ink and White Opaque body imprinted with 30 in black ink filled with White to Off-white powder.  
• Capsules 40 mg: Hard Gelatin Capsule Shell Size "3" Blue Green Opaque Cap imprinted with AC in Black ink and White Opaque body imprinted with 40 in black ink filled with White to Off-white powder.  
• Capsules 50 mg: Hard Gelatin Capsule Shell Size "3" Blue Opaque Cap imprinted with AC in Black ink and White Opaque body imprinted with 50 in black ink filled with White to Off-white powder.  
• Capsules 60 mg: Hard Gelatin Capsule Shell Size "2" Aqua Blue Opaque Cap imprinted with AC in Black ink and Aqua Blue Opaque body imprinted with 60 in black ink filled with White to Off-white powder.  
• Capsules 70 mg: Hard Gelatin Capsule Shell Size "2" White Opaque Cap imprinted with AC in Black ink and Blue Transparent body imprinted with 70 in black ink filled with White to Off-white powder.

4 CONTRAINDICATIONS

Lisdexamfetamine dimesylate capsules are contraindicated in patients with:  
• Known hypersensitivity to amphetamine products or other ingredients in lisdexamfetamine dimesylate capsules. Anaphylactic reactions, Stevens-Johnson Syndrome, angioedema, and urticaria have been observed in postmarketing reports (See *Adverse Reactions (6.2)*).

• Patients taking monoamine oxidase inhibitors (MAOIs) or within 14 days of stopping MAOIs including such as linezolid or intravenous methylene blue, because of an increased risk of hypertensive crisis (See **Warnings and Precautions (5.7)** and *Drug Interactions (7.1)*).

5 WARNINGS AND PRECAUTIONS

5.1 Abuse, Misuse, and Addiction

Lisdexamfetamine dimesylate capsules has a high potential for abuse and misuse. The use of lisdexamfetamine dimesylate capsules exposes individuals to the risks of abuse and misuse, which can lead to the development of a substance use disorder, including addiction. Lisdexamfetamine dimesylate capsules can be diverted for non-medical use into illicit channels or distribution (See *Drug Abuse and Dependence (8.2)*). Misuse and abuse of CNS stimulants, including lisdexamfetamine dimesylate capsules, can result in overdose and death (See *Overdose (10)*), and this risk is increased with higher doses or unapproved methods of administration, such as snorting or injection.

Before prescribing lisdexamfetamine dimesylate capsules, assess each patient's risk for abuse, misuse, and addiction. Educate patients and their families about these risks and proper disposal of any unused drug. Advise patients to store lisdexamfetamine dimesylate capsules in a safe place, preferably locked, and instruct patients to not give lisdexamfetamine dimesylate capsules to anyone else. Throughout lisdexamfetamine dimesylate capsules treatment, reassess each patient's risk of abuse, misuse, and addiction and frequently monitor for signs and symptoms of abuse, misuse, and addiction.

5.2 Risks to Patients with Serious Cardiac Disease

Sudden death has been reported in patients with structural cardiac abnormalities or other serious cardiac disease who were treated with CNS stimulants at the recommended ADHD dosage. Avoid lisdexamfetamine dimesylate capsules use in patients with known structural cardiac abnormalities, cardiomyopathy, serious cardiac arrhythmia, coronary artery disease, or other serious cardiac disease.

5.3 Increased Blood Pressure and Heart Rate

CNS stimulants cause an increase in blood pressure (mean increase about 2 to 4 mm Hg) and heart rate (mean increase about 3 to 5 bpm). Some patients may have larger increases.

Monitor all lisdexamfetamine dimesylate capsules-treated patients for potential tachycardia and hypertension.

5.4 Psychiatric Adverse Reactions

Exacerbation of Pre-existing Psychosis

CNS stimulants may exacerbate symptoms of behavior disturbance and thought disorder in patients with a pre-existing psychotic disorder.

Induction of a Manic Episode in Patients with Bipolar Disorder  
CNS stimulants may induce a manic or mixed episode. Prior to initiating lisdexamfetamine dimesylate capsules treatment, screen patients for risk factors for developing a manic episode (e.g., comorbid or history of depressive symptoms or a family history of suicide, bipolar disorder, and depression).

New Psychotic or Manic Symptoms

CNS stimulants, at the recommended dosage, may cause psychotic or manic symptoms (e.g., hallucinations, delusional thought, or mania) in patients without a prior history of psychotic illness or mania. In a pooled analysis of multiple short-term, placebo-controlled studies of CNS stimulants conducted in approximately 0.1% of CNS stimulant-treated patients, psychotic symptoms compared to 0% of placebo-treated patients. If such symptoms occur, consider discontinuing lisdexamfetamine dimesylate capsules.

5.5 Long-Term Suppression of Growth in Pediatric Patients

Lisdexamfetamine dimesylate capsules are not approved for use and are not recommended in pediatric patients below 6 years of age (See *Use in Specific Populations (8.4)*).

In 4-week, placebo-controlled trial of lisdexamfetamine dimesylate capsules in pediatric patients ages 6 to 12 years old with ADHD, there was a dose-related decrease in weight in the lisdexamfetamine dimesylate capsules groups compared to weight gain in the placebo group. Additionally, in another study, there was slowing of the increase in height (See *Adverse Reactions (6.1)*).

Closely monitor growth weight and height in lisdexamfetamine dimesylate capsules-treated pediatric patients. Patients who are not growing or gaining height or weight as expected may need to have their treatment interrupted.

5.6 Peripheral Vascopathy, including Raynaud's Phenomenon

CNS stimulants, including lisdexamfetamine dimesylate capsules, used to treat ADHD are associated with peripheral vascopathy, including Raynaud's phenomenon. Signs and symptoms are usually intermittent and mild; however, sequelae have included digital ulceration and/or self tissue breakdown. Effects of peripheral vascopathy, including Raynaud's phenomenon, were observed in post-marketing reports and at the therapeutic dosages of CNS stimulants in all age groups throughout the course of treatment.

- End stage renal disease (ESRD): Maximum dose is 30 mg/day (2,5)

DOSEAGE FORMS AND STRENGTHS

- Capsules: 10 mg, 20 mg, 30 mg, 40 mg, 50 mg, 60 mg, 70 mg (3)

CONTRAINDICATIONS

- Known hypersensitivity to amphetamine products or other ingredients in lisdexamfetamine dimesylate capsules (4)
- Use with monoamine oxidase (MAO) inhibitor or within 14 days of the last MAO inhibitor dose (4, 7.1)

WARNINGS AND PRECAUTIONS

- Risks to Patients with Serious Cardiac Disease: Avoid use in patients with known structural cardiac abnormalities, cardiomyopathy, serious cardiac arrhythmia, coronary artery disease, or other serious cardiac disease (5.2)
- Increased Blood Pressure and Heart Rate: Monitor blood pressure and pulse. (5.3)
- Psychiatric Adverse Reactions: Prior to initiating lisdexamfetamine dimesylate capsules, screen patients for risk factors for developing a manic episode. If new psychotic or manic symptoms occur, consider discontinuing lisdexamfetamine dimesylate capsules. (5.4)

- Long-Term Suppression of Growth in Pediatric Patients: Closely monitor growth (height and weight) in pediatric patients. Lisdexamfetamine is not growing or gaining height or weight as expected may need to have their treatment interrupted. (5.5)
- Peripheral Vascopathy, including Raynaud's phenomenon: Careful observation for digital changes is necessary during lisdexamfetamine dimesylate capsules treatment. Further clinical evaluation (e.g., rheumatology referral) may be appropriate for patients who develop signs or symptoms of peripheral vascopathy. (5.6)

- Serotonin Syndrome: Increased risk when co-administered with serotonergic agents (e.g., SSRIs, SNRIs, triptans), but also during overdose situations. If or occurs, discontinue lisdexamfetamine dimesylate capsules and initiate supportive treatment. (4, 5.7, 10)
- Motor and Verbal Tics, and Worsening of Tourette's Syndrome: Before initiating lisdexamfetamine dimesylate capsules, assess the family history and clinically evaluate patients for tics or Tourette's syndrome. Regularly monitor patients for the emergence or worsening of tics or Tourette's syndrome. Discontinue treatment if clinically appropriate. (5.8)

ADVERSE REACTIONS

Most common adverse reactions (incidence >5% and at a rate at least twice placebo) in pediatric patients ages 6 to 17 years, and/or adults with ADHD were: anorexia, anxiety, decreased appetite, decreased weight, diarrhea, dizziness, dry mouth, irritability, insomnia, nausea, upper abdominal pain, and vomiting. (6.1)

Most common adverse reactions (incidence >5% and at a rate at least twice placebo) in adults with BED were dry mouth, insomnia, decreased weight, decreased appetite, decreased heart rate, and decreased blood pressure. (6.2)

To report SUSPECTED ADVERSE REACTIONS, contact Carter Pharmaceuticals, Inc., at 1-866-495-4330 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

• **DRUG ABUSE AND DEPENDENCE**  
Lisdexamfetamine dimesylate capsules have a high potential for abuse and misuse, which can lead to the development of a substance use disorder, including addiction. Misuse and abuse of CNS stimulants, including lisdexamfetamine dimesylate capsules, can result in overdose and death (5.1, 8.2, 10).

USE IN SPECIFIC POPULATIONS

- Pregnancy may cause fetal harm (8.1)
- Lactation: Breastfeeding not recommended (8.2)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 10/2025

|                                                      |                                                               |
|------------------------------------------------------|---------------------------------------------------------------|
| 8.2 Lactation                                        | 8.4 Pediatric Use                                             |
| 8.5 Geriatric Use                                    | 8.6 Renal Impairment                                          |
| 9.1 Controlled Substance                             | 9.2 Abuse                                                     |
| 9.3 Dependence                                       | 9.4 Dependence                                                |
| 10 OVERDOSE                                          | 10.1 Description                                              |
| 11 DESCRIPTION                                       | 11.1 Clinical Pharmacology                                    |
| 12.1 Mechanism of Action                             | 12.2 Pharmacokinetics                                         |
| 12.3 Pharmacokinetics                                | 12.4 Pharmacokinetics                                         |
| 13 NONCLINICAL TOXICOLOGY                            | 13.1 Carcinogenesis, Mutagenesis, and Impairment of Fertility |
| 13.2 Animal Toxicology and/or Pharmacology           | 14 CLINICAL STUDIES                                           |
| 14.1 Attention Deficit Hyperactivity Disorder (ADHD) | 14.2 Binge Eating Disorder (BED)                              |
| 16 HOW SUPPLIED/STORAGE AND HANDLING                 | 16.1 How Supplied                                             |
| 16.2 Storage and Handling                            | 17 PATIENT COUNSELING INFORMATION                             |
| 17.1 Patient Counseling Information                  | 17.2 Patient Counseling Information                           |

\*Sections or subsections omitted from the full prescribing information are not listed.

Signs and symptoms generally improved after dosage reduction or discontinuation of the CNS stimulant.

Careful observation for digital changes is necessary during lisdexamfetamine dimesylate capsules treatment. Further clinical evaluation (e.g., rheumatology referral) may be appropriate for lisdexamfetamine dimesylate capsules-treated patients who develop signs or symptoms of peripheral vascopathy.

5.7 Serotonin Syndrome

Lisdexamfetamine, a potentially life-threatening reaction, may occur when amphetamines are used in combination with other drugs that affect the serotonergic neurotransmitter systems such as monoamine oxidase inhibitors (MAOIs), selective serotonin reuptake inhibitors (SSRIs), serotonin norepinephrine reuptake inhibitors (SNRIs), triptans, tricyclic antidepressants, fentanyl, lithium, tramadol, typhoid fever drugs (e.g., fluoroquinolones), and other serotonergic agents (e.g., SSRIs, SNRIs, triptans), but also during overdose situations. If or occurs, discontinue lisdexamfetamine dimesylate capsules and initiate supportive treatment. (4, 5.7, 10)

Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, delirium, and coma), autonomic instability (e.g., tachycardia, labile blood pressure, dizziness, diaphoresis, flushing, hyperreflexia), neuromuscular symptoms (e.g., tremor, rigidity), myoclonus, hyperreflexia, incoordination, seizures, and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhea).

Concomitant use of lisdexamfetamine dimesylate capsules with MAOIs is contraindicated (See **Contraindications (4)**).

Discontinue treatment with lisdexamfetamine dimesylate capsules and any concomitant serotonergic agents immediately if symptoms of serotonin syndrome occur. Initiate supportive symptomatic treatment. If concomitant use of lisdexamfetamine dimesylate capsules with other serotonergic drugs or CYP2D6 inhibitors is clinically warranted, initiate lisdexamfetamine dimesylate capsules with lower doses, monitor patients for the emergence of serotonin syndrome during drug initiation or titration, and inform patients of the increased risk for serotonin syndrome. (See **Warnings and Precautions (5.7)** and *Drug Interactions (7.1)*).

**5.8 Motor and Verbal Tics, and Worsening of Tourette's Syndrome**  
CNS stimulants, including amphetamines, have been associated with the onset or exacerbation of motor and verbal tics. Worsening of Tourette's syndrome has also been reported (See *Adverse Reactions (6.2)*).

Before initiating lisdexamfetamine dimesylate capsules, assess the family history and clinically evaluate patients for tics or Tourette's syndrome. Regularly monitor lisdexamfetamine dimesylate capsules-treated patients for the emergence or worsening of tics or Tourette's syndrome, and discontinue treatment if clinically appropriate.

6 ADVERSE REACTIONS

The following adverse reactions are discussed in greater detail in other sections of the labeling:  
• Known hypersensitivity to amphetamine products or other ingredients in lisdexamfetamine dimesylate capsules (See **Contraindications (4)**)

• Hypertensive Crisis When Used Concomitantly with Monoamine Oxidase Inhibitors (See **Contraindications (4)** and *Drug Interactions (7.1)*)

• Abuse, Misuse, and Addiction (See **Boxed Warning, Warnings and Precautions (5.1)**, and *Drug Abuse and Dependence (8.2, 9, 3)*)

• Risks to Patients with Serious Cardiac Disease (See **Warnings and Precautions (5.2)**)

• Increased Blood Pressure and Heart Rate (See **Warnings and Precautions (5.3)**)

• Psychiatric Adverse Reactions (See **Warnings and Precautions (5.4)**)

• Long-Term Suppression of Growth in Pediatric Patients (See **Warnings and Precautions (5.5)**)

• Peripheral Vascopathy, including Raynaud's phenomenon (See **Warnings and Precautions (5.6)**)

• Serotonin Syndrome (See **Warnings and Precautions (5.7)**)

• Motor and Verbal Tics, and Worsening of Tourette's Syndrome (See **Warnings and Precautions (5.8)**)

6.1 Clinical Trials Experience

Clinical trials are conducted under widely varying conditions; adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice. Attention Deficit Hyperactivity Disorder.

The safety data in this section is based on data from the 4-week controlled parallel-group clinical studies of lisdexamfetamine dimesylate capsules in pediatric and adult patients with ADHD (See *Clinical Studies (14.1)*).

**Adverse Reactions Associated with Discontinuation of Treatment in ADHD Clinical Trials**  
In the controlled trial in pediatric patients ages 6 to 12 years (Study 1), 8% (18/217) of lisdexamfetamine dimesylate capsules-treated patients discontinued due to adverse reactions compared to 0% (0/72) of placebo-treated patients. The most frequently reported adverse reactions (1% or more and twice the rate of placebo) were ECG voltage criteria for ventricular hypertrophy, tic, dry mouth, psychomotor hyperactivity, insomnia, decreased appetite and rash (1 instance each adverse reaction, i.e., >2/18 (11%) less frequently reported adverse reactions (less than 1% or less than twice the rate of placebo) included abdominal pain (upper), dry mouth, weight decreased, dizziness, somnolence, lightheadedness, chest pain, anger, and hypertension.

In the controlled trial in pediatric patients ages 13 to 17 years (Study 4), 3% (23/733) of lisdexamfetamine dimesylate capsules-treated patients discontinued due to adverse reactions compared to 1% (1/77) of placebo-treated patients. The most frequently reported adverse reactions (1% or more and twice the rate of placebo) were decreased appetite (2/233; 1%) and insomnia (2/233; 1%). Less frequently reported adverse reactions (less than 1% or less than twice the rate of placebo) included irritability, dermatitis, anxiety, and depression.

In the controlled adult trial (Study 7), 6% (21/358) of lisdexamfetamine dimesylate capsules-treated patients discontinued due to adverse reactions compared to 2% (1/62) of placebo-treated patients. The most frequently reported adverse reactions (1% or more and twice the rate of placebo) were insomnia (8/358; 2%), tachycardia (3/358; 1%), hypertension (3/358; 1%), headache (2/358; 1%), anxiety (2/358; 1%), and dyspnea (3/358; 1%). Less frequently reported adverse reactions (less than 1% or less than twice the rate of placebo) included palpitations, diarrhea, nausea, decreased appetite, dizziness, agitation, depression, and irritability.

**Adverse Reactions Occurring at an Incidence of >5% or More Among Lisdexamfetamine Dimesylate Capsules Treated Patients with ADHD in Clinical Trials**  
The most common adverse reactions (incidence >5% and at a rate at least twice placebo) reported in pediatric patients ages 6 to 17 years, and/or adults were anorexia, anxiety, decreased appetite, decreased weight, diarrhea, dizziness, dry mouth, irritability, insomnia, nausea, upper abdominal pain, and vomiting.

**Adverse Reactions Occurring at an Incidence of 2% or More Among Lisdexamfetamine Dimesylate Capsules Treated Patients with ADHD in Clinical Trials**  
In the controlled trial in pediatric patients ages 6 to 12 years (Study 1), 6% (23/358) of lisdexamfetamine dimesylate capsules-treated patients discontinued due to adverse reactions compared to 2% (1/62) of placebo-treated patients. The most frequently reported adverse reactions (1% or more and twice the rate of placebo) were insomnia (8/358; 2%), tachycardia (3/358; 1%), hypertension (3/358; 1%), headache (2/358; 1%), anxiety (2/358; 1%), and dyspnea (3/358; 1%). Less frequently reported adverse reactions (less than 1% or less than twice the rate of placebo) included palpitations, diarrhea, nausea, decreased appetite, dizziness, agitation, depression, and irritability.

**Table 1 Adverse Reactions Reported by 2% or More of Pediatric Patients Ages 6 to 12 Years with ADHD Taking Lisdexamfetamine Dimesylate Capsules and Greater than or Equal to Twice the Incidence in Patients Taking Placebo in a 4-Week Clinical Trial (Study 1)**

| Lisdexamfetamine dimesylate capsules (n=217) | Placebo (n=72) |
|----------------------------------------------|----------------|
| Decreased Appetite                           | 3%             |
| Anxiety                                      | 4%             |
| Abdominal Pain Upper                         | 2%             |
| Insomnia                                     | 1%             |
| Vomiting                                     | 4%             |
| Weight Decreased                             | 1%             |
| Nausea                                       | 3%             |
| Dry Mouth                                    | 3%             |
| Dizziness                                    | 0%             |
| Affectability                                | 0%             |
| Rash                                         | 0%             |
| Pyrexia                                      | 0%             |
| Somnolence                                   | 1%             |
| Tic                                          | 2%             |
| Anorexia                                     | 0%             |

**Table 2 Adverse Reactions Reported by 2% or More of Pediatric Patients Ages 13 to 17 Years with ADHD Taking Lisdexamfetamine Dimesylate Capsules and Greater than or Equal to Twice the Incidence in Patients Taking Placebo in a 4-Week Clinical Trial (Study 4)**

| Lisdexamfetamine dimesylate capsules (n=233) | Placebo (n=77) |
|----------------------------------------------|----------------|
| Decreased Appetite                           | 2%             |
| Insomnia                                     | 2%             |
| Weight Decreased                             | 0%             |
| Dry Mouth                                    | 1%             |
| Palpitations                                 | 1%             |
| Anorexia                                     | 2%             |
| Tremor                                       | 0%             |

**Table 3 Adverse Reactions Reported by 2% or More of Adult Patients with ADHD Taking Lisdexamfetamine Dimesylate Capsules and Greater than or Equal to Twice the Incidence in Patients Taking Placebo in a 4-Week Clinical Trial (Study 7)**

| Lisdexamfetamine dimesylate capsules (n=58) | Placebo (n=62) |
|---------------------------------------------|----------------|
| Decreased Appetite                          | 2%             |