

MEDICATION GUIDE

Eltrombopag

(el-TROM-boe-pag)

Tablets

What is the most important information I should know about eltrombopag tablets?

Eltrombopag tablets can cause serious side effects, including:

Liver problems:

- If you have chronic hepatitis C virus and take eltrombopag tablets with interferon and ribavirin treatment, eltrombopag tablets may increase your risk of liver problems. If your healthcare provider tells you to stop your treatment with interferon and ribavirin, you will also need to stop taking eltrombopag tablets.
- Eltrombopag tablets may increase your risk of liver problems that may be severe and possibly life threatening. Your healthcare provider will do blood tests to check your liver function before you start taking eltrombopag tablets and during your treatment. Your healthcare provider may stop your treatment with eltrombopag tablets if you have changes in your liver function blood tests.

Tell your healthcare provider right away if you have any of these signs and symptoms of liver problems:

- yellowing of the skin or the whites of the eyes (jaundice)
- unusual darkening of the urine
- unusual tiredness

- right upper stomach area (abdomen) pain
- confusion
- swelling of the stomach area (abdomen)

See “What are the possible side effects of eltrombopag tablets?” for other side effects of eltrombopag tablets.

What are eltrombopag tablets?

Eltrombopag tablet is a prescription medicine used to treat adults and children 1 year of age and older with low blood platelet counts due to persistent or chronic immune thrombocytopenia (ITP), when other medicines to treat ITP or surgery to remove the spleen have not worked well enough.

Eltrombopag tablet are also used to treat people with:

- low blood platelet counts due to chronic hepatitis C virus (HCV) infection before and during treatment with interferon.
- severe aplastic anemia (SAA) when other medicines to treat SAA have not worked well enough.

Eltrombopag tablets are used to try to raise platelet counts in order to lower your risk for bleeding.

Eltrombopag tablets are not used to make platelet counts normal.

Eltrombopag tablets are not for use in people with a pre-cancerous condition called myelodysplastic syndrome (MDS), or in people with low platelet counts caused by certain other medical conditions or diseases.

It is not known if eltrombopag tablets are safe and effective when used with other antiviral medicines to treat chronic hepatitis C.

It is not known if eltrombopag tablets are safe and effective in children:

- younger than 1 year with ITP
- with low blood platelet counts due to chronic hepatitis C
- whose severe aplastic anemia (SAA) has not improved after previous treatments
- younger than 2 years when used in combination with other medicines to treat SAA as the first treatment for SAA

Before you take eltrombopag tablets, tell your healthcare provider about all your medical conditions, including if you:

- have liver problems
- have a precancerous condition called MDS or a blood cancer
- have or had a blood clot
- have a history of cataracts
- have had surgery to remove your spleen (splenectomy)
- have bleeding problems
- are of East-/Southeast-Asian ancestry. You may need a lower dose of eltrombopag tablets
- are pregnant or plan to become pregnant. It is not known if eltrombopag tablets will harm an unborn baby. Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with eltrombopag tablets.
  - Females who are able to become pregnant, should use effective birth control (contraception) during treatment with eltrombopag tablets and for at least 7 days after stopping treatment with eltrombopag tablets. Talk to your healthcare provider about birth control methods that may be right for you during this time.
- are breastfeeding or plan to breastfeed. You should not breastfeed during your treatment with eltrombopag tablets. Talk to your healthcare provider about the best way to feed your baby during this time.

- Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Eltrombopag tablets may affect the way certain medicines work. Certain other medicines may affect the way eltrombopag tablets works.

Especially tell your healthcare provider if you take:

- certain medicines used to treat high cholesterol, called “statins”
- a blood thinner medicine

Certain medicines may keep eltrombopag tablets from working correctly. Take eltrombopag tablets at least 2 hours before or 4 hours after taking these products:

- antacid medicine used to treat stomach ulcers or heartburn
- multivitamins or products that contain iron, calcium, aluminum, magnesium, selenium, and zinc which may be found in mineral supplements

Ask your healthcare provider if you are not sure if your medicine is one that is listed above.


Know the medicines you take. Keep a list of them and show it to your healthcare provider and pharmacist when you get a new medicine.

How should I take eltrombopag tablets?

- Take eltrombopag tablets exactly as your healthcare provider tells you to take them. Your healthcare provider will prescribe the dose of eltrombopag tablets that is right for you.
- If your healthcare provider prescribes eltrombopag tablets, take eltrombopag tablets whole.

Do not split, chew, or crush eltrombopag tablets and do not mix with food or liquids.

- Do not** stop taking eltrombopag tablets without talking with your healthcare provider first. Do not change your dose or schedule for taking eltrombopag tablets unless your healthcare provider tells you to change them.
- Take eltrombopag tablets without a meal or with a meal low in calcium (50 mg or less) and at least 2 hours before or 4 hours after eating calcium-rich foods, such as dairy products, calcium-fortified juices, and certain fruits and vegetables.
- If you miss a dose of eltrombopag, wait and take your next scheduled dose. Do not take more than 1 dose of eltrombopag tablets in 1 day.

Artwork information			
Customer	Comber	Market	USA
Dimensions (mm)	200 x 350 mm	Non Printing Colors	 Die cut
Pharma Code No.	NA		
Printing Colours (01)	Black		
Others: Pharma code position and Orientation are tentative, will be changed based on folding size.			

