

Unvarnish Area For Batch Coding

21002723



Rx only

NDC 31722-224-31

Voriconazole for Injection

200 mg* per vial

For Intravenous Infusion Only

Sterile Single-Dose Vial

Discard Unused Portion

1 Vial

DOSAGE AND USE:

Reconstitute with 19 mL of Water for Injection to give a clear solution containing 10 mg/mL voriconazole and an extractable volume of 20 mL. Must be further diluted before use. For administration, the required volume of the reconstituted solution is added to a recommended compatible infusion solution to provide a final voriconazole solution containing 0.5 to 5 mg/mL. For appropriate diluents and storage recommendations, refer to prescribing information.

FOR INTRAVENOUS INFUSION AFTER DILUTION.

* With reconstitution each mL contains 10 mg voriconazole and 160 mg sulfobutyl ether β-cyclodextrin sodium (SBECD).

Store at 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].



Rx only

NDC 31722-224-31

Voriconazole for Injection

200 mg* per vial

For Intravenous Infusion Only

Sterile Single-Dose Vial

Discard Unused Portion

1 Vial

The vial stoppers are not made with natural rubber latex.

No preservatives

U.S. Contact Number: 1-866-495-1995

Manufactured for: Camber Pharmaceuticals, Inc. Piscataway, NJ 08854.

Manufactured by:



ASPIRO PHARMA LIMITED

Survey No. 321, Biotech Park, Phase-III, Karkapatla Village, Markook Mandal, Siddipet Dist., Telangana State - 502281, INDIA.

Mfg. Lic. No.: 36/MD/AP/2013/F/G



Dimensions: 42x 42 x 90 mm

PANTONE 2736 C

PANTONE 1805 C

Black