

2D Data Matrix to be printed with serial number on each leaflet. The number should not be repeated

Note: Pharmacode position and orientation will be changed as per folding dimensions

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Nilotinib CAPSULES safely and effectively. See full prescribing information for Nilotinib CAPSULES.

NILOTINIB CAPSULES, for oral use
Initial U.S. Approval: 2007

WARNING: QTc Prolongation and Sudden Deaths

See full prescribing information for complete boxed warning.

- Nilotinib prolongs the QT interval. Prior to nilotinib administration and periodically, monitor for hypokalemia or hypomagnesemia and correct deficiencies. (5.2) Obtain ECGs to monitor the QTc at baseline, seven days after initiation, and periodically thereafter, and following any dose adjustments (2, 5.3, 5.7, 5.12).
- Sudden deaths have been reported in patients receiving nilotinib. (5.3) Do not administer nilotinib to patients with hypokalemia, hypomagnesemia, or long QT syndrome. (4, 5.2)
- Avoid use of concomitant drugs known to prolong the QT interval and strong CYP3A4 inhibitors. (7.1, 7.2)
- Avoid food 2 hours before and 1 hour after taking the dose. (2.1)

INDICATIONS AND USAGE

- Adult patients with newly diagnosed Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in chronic phase (1.1)
- Adult patients with chronic phase (CP) and accelerated phase (AP) Ph+ CML resistant to or intolerant to prior therapy that included imatinib. (1.2)

DOSEAGE AND ADMINISTRATION

- Recommended Adult Dose: Newly diagnosed Ph+ CML-CP: 300 mg orally twice daily. Resistant or intolerant Ph+ CML-CP and CML-AP: 400 mg orally twice daily. (2.1)
- See Dosage and Administration for full dosing instructions and dose- reduction instructions for toxicity. (2.1)
- Reduce starting dose in patients with baseline hepatic impairment. (2.7)
- Eligible newly diagnosed adult patients with Ph+ CML-CP who have received nilotinib capsules for a minimum of 3 years and have achieved a sustained molecular response (MR4.5) and patients with Ph+ CML-CP resistant or intolerant to imatinib who have received nilotinib capsules for at least 3 years and have achieved a sustained molecular response (MR4.5) may be considered for treatment discontinuation. (2.2, 2.3, 5.16)

DOSEAGE FORMS AND STRENGTHS

Capsules: 150 mg, and 200 mg (3)

CONTRAINDICATIONS

Nilotinib capsules are contraindicated in patients with hypokalemia, hypomagnesemia, or long QT syndrome. (4)

WARNINGS AND PRECAUTIONS

- Myelosuppression: Monitor complete blood count (CBC) during therapy and manage by treatment interruption or dose reduction. (5.1)

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: QTc Prolongation and Sudden Deaths

1. INDICATIONS AND USAGE
 - 1.1 Adult Patients With Newly Diagnosed Ph+ CML-CP
 - 1.2 Adult Patients With Resistant or Intolerant Ph+ CML-CP and CML-AP
2. DOSAGE AND ADMINISTRATION
 - 2.1 Recommended Dose
 - 2.2 Discontinuation of Treatment After a Sustained Molecular Response (MR4.5) on Nilotinib Capsules
 - 2.3 Reinitiation of Treatment in Patients Who Lose Molecular Response After Discontinuation of Therapy With Nilotinib Capsules
 - 2.4 Dosage Modification for QT Interval Prolongation
 - 2.5 Dosage Modifications for Myelosuppression
 - 2.6 Dosage Modifications for Selected Non-Hematologic Laboratory Abnormalities and Other Toxicities
 - 2.7 Dosage Modification for Hepatic Impairment
 - 2.8 Dosage Modification With Concomitant Strong CYP3A4 Inhibitors
3. DOSAGE FORMS AND STRENGTHS
4. CONTRAINDICATIONS
5. WARNINGS AND PRECAUTIONS
 - 5.1 Myelosuppression
 - 5.2 QT Prolongation
 - 5.3 Sudden Deaths
 - 5.4 Cardiac and Arterial Vascular Occlusive Events
 - 5.5 Pancreatitis and Elevated Serum Lipase
 - 5.6 Hepatotoxicity
 - 5.7 Electrolyte Abnormalities
 - 5.8 Tumor Lysis Syndrome
 - 5.9 Hemorrhage
 - 5.10 Total Gastroctomy
 - 5.11 Monitoring Laboratory Tests
 - 5.12 Fluid Retention
 - 5.13 Effects on Growth and Development in Pediatric Patients
 - 5.14 Embryo-Fetal Toxicity

FULL PRESCRIBING INFORMATION

WARNING: QTc Prolongation and Sudden Deaths

- Nilotinib prolongs the QT interval. Prior to nilotinib administration and periodically, monitor for hypokalemia or hypomagnesemia and correct deficiencies. (See Warnings and Precautions (5.2)).
- Obtain ECGs to monitor the QTc at baseline, seven days after initiation, and periodically thereafter, and following any dose adjustments. (See Warnings and Precautions (5.2, 5.3, 5.7, 5.12)).
- Sudden deaths have been reported in patients receiving nilotinib. (See Warnings and Precautions (5.3)).
- Do not administer nilotinib to patients with hypokalemia, hypomagnesemia, or long QT syndrome. (See Contraindications (4), Warnings and Precautions (5.2)).
- Avoid use of concomitant drugs known to prolong the QT interval and strong CYP3A4 inhibitors. (See Drug Interactions (7.1, 7.2)).
- Avoid food 2 hours before and 1 hour after taking the dose. (See Dosage and Administration (2.1)).

1. INDICATIONS AND USAGE

1.1 Adult Patients With Newly Diagnosed Ph+ CML-CP
Nilotinib capsules are indicated for the treatment of adult patients with newly diagnosed Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in chronic phase.

Additional pediatric use information is approved for Novartis Pharmaceuticals Corporation's TASIGNA (nilotinib) capsules. However, due to Novartis Pharmaceuticals Corporation's marketing exclusivity rights, this drug product is not labeled with that pediatric information.

1.2 Adult Patients With Resistant or Intolerant Ph+ CML-CP and CML-AP
Nilotinib capsules are indicated for the treatment of adult patients with chronic phase and accelerated phase Philadelphia chromosome positive chronic myelogenous leukemia (Ph+ CML) resistant or intolerant to prior therapy that included imatinib.

Additional pediatric use information is approved for Novartis Pharmaceuticals Corporation's TASIGNA (nilotinib) capsules. However, due to Novartis Pharmaceuticals Corporation's marketing exclusivity rights, this drug product is not labeled with that pediatric information.

2. DOSAGE AND ADMINISTRATION

2.1 Recommended Dose

Dose nilotinib capsules twice daily at approximately 12-hour intervals on an empty stomach. No food should be consumed for at least 2 hours before the dose is taken and for at least 1 hour after the dose is taken. Advise patients to swallow the capsules whole with water. (See Boxed Warning, Clinical Pharmacology (12.3)).

For patients who are unable to swallow capsules, the contents of each capsule may be dispersed in 1 teaspoon of applesauce (purged apple). The mixture should be taken immediately (within 15 minutes) and should not be stored for future use. (See Clinical Pharmacology (12.3)).

Additional pediatric use information is approved for Novartis Pharmaceuticals Corporation's TASIGNA (nilotinib) capsules. However, due to Novartis Pharmaceuticals Corporation's marketing exclusivity rights, this drug product is not labeled with that pediatric information.

2.2 Discontinuation of Treatment After a Sustained Molecular Response (MR4.5) on Nilotinib Capsules
Eligibility for discontinuation of treatment.

Ph+ CML-CP patients with typical BCR-ABL transcripts, who have been taking nilotinib capsules for a minimum of 3 years and have achieved a sustained molecular response (MR4.5, corresponding to = BCR-ABL/ABL $\leq 0.0032\%$ IS), may be eligible for treatment discontinuation. (See Clinical Studies (14.3, 14.4)). Information on the detection and quantitation of BCR-ABL transcripts in patients with chronic phase disease to determine eligibility for treatment discontinuation is available at <http://www.fda.gov/compliance/patient-safety>.

Patients with typical BCR-ABL transcripts (e13a2/b2a2 or e142b2/b3a2), who achieve the sustained MR4.5 criteria, are eligible for discontinuation of nilotinib capsules. Patients must continue to be monitored for possible loss of molecular remission after treatment discontinuation. Use the same FDA-authorized test to consistently monitor molecular response levels while on and off treatment. Consider discontinuation in patients with Ph+ CML-CP who have:

- been treated with nilotinib capsules for at least 3 years
- maintained a molecular response of at least MR4.0 (corresponding to = BCR-ABL/ABL $\leq 0.01\%$ IS) for one year prior to discontinuation of therapy
- achieved an MR4.5 for the last assessment taken immediately prior to discontinuation of therapy
- been confirmed to express the typical BCR-ABL transcripts (e13a2/b2a2 or e142b2/b3a2)
- no history of accelerated phase or blast crisis
- no history of prior attempts of treatment

Consider discontinuation in patients with Ph+ CML-CP that are resistant or intolerant to imatinib who have achieved a sustained molecular response (MR4.5) on nilotinib capsules who have:

- been treated with nilotinib capsules for a minimum of 3 years
- been treated with imatinib only prior to treatment with nilotinib capsules
- achieved a molecular response of MR4.5 (corresponding to = BCR-ABL/ABL $\leq 0.0032\%$ IS)
- sustained an MR4.5 for a minimum of one year immediately prior to discontinuation of therapy
- been confirmed to express the typical BCR-ABL transcripts (e13a2/b2a2 or e142b2/b3a2)
- no history of accelerated phase or blast crisis
- no history of prior attempts of treatment

Monitor BCR-ABL transcript levels and complete blood count (CBC) with differential in patients who have discontinued nilotinib capsules therapy monthly for one year, then every 6 weeks for the second year, and every 12 weeks thereafter. (See Warnings and Precautions (5.16)).

Upon the loss of MR4.0 (corresponding to = BCR-ABL/ABL $\leq 0.01\%$ IS) during the treatment-free phase, monitor BCR-ABL transcript levels every 2 weeks until BCR-ABL levels remain lower than major molecular response (MMR), corresponding to MR5.0 (= BCR-ABL/ABL $\leq 0.1\%$ IS) for 4 consecutive measurements. The patient can then proceed to the original monitoring schedule.

2.3 Reinitiation of Treatment in Patients Who Lose Molecular Response After Discontinuation of Therapy With Nilotinib Capsules

- Newly diagnosed patients who lose MMR must reinitiate treatment within 4 weeks at the dose level prior to discontinuation of therapy. (See Warnings and Precautions (5.16)). Patients who reinitiate nilotinib capsules therapy should have their BCR-ABL transcript levels monitored monthly until major molecular response is re-established and every 12 weeks thereafter.
- Patients resistant or intolerant to prior treatment that included imatinib with confirmed loss of MR4.0 (2 consecutive measures separated by at least 4 weeks showing loss of MR4.0) or loss of MMR must reinitiate treatment within 4 weeks at the dose level prior to discontinuation of therapy. (See Warnings and Precautions (5.16)). Patients who reinitiate nilotinib capsules therapy should have their BCR-ABL transcript levels monitored monthly until previous major molecular response or MR4.0 is re-established and every 12 weeks thereafter.

2.4 Dosage Modification for QT Interval Prolongation
See Table 2 for dose adjustments for QT interval prolongation. (See Warnings and Precautions (5.2), Clinical Pharmacology (12.2)).

Table 2: Dosage Adjustments for Adult Patients With QT Prolongation

Degree of QTc prolongation	Dosage adjustment
ECGs with a QTc greater than 480 msec	1. Withhold nilotinib capsules, and perform an analysis of serum potassium greater than 4.0 mmol/L and magnesium, and if below lower limit of normal, correct with supplements to return within normal limits. Concomitant medication use must be reviewed. Resume within 2 weeks at prior dose if QTc returns to less than 450 msec, and to within 20 msec of baseline.
	3. If QTc is between 450 msec and 480 msec after 2 weeks, reduce the dose to 400 mg once daily in adult patients.
	4. Discontinue nilotinib capsules if, following dose-reduction to 400 mg once daily in adult patients, QTc returns to greater than 480 msec.
	5. An ECG should be repeated approximately 7 days after any dose adjustment.

Abbreviation: ECG, electrocardiogram.

Additional pediatric use information is approved for Novartis Pharmaceuticals Corporation's TASIGNA (nilotinib) capsules. However, due to Novartis Pharmaceuticals Corporation's marketing exclusivity rights, this drug product is not labeled with that pediatric information.

2.5 Dosage Modifications for Myelosuppression
Withhold or reduce nilotinib capsules dosage for hematological toxicities (neutropenia, thrombocytopenia) that are not related to underlying leukemia (Table 3). (See Warnings and Precautions (5.1)).

- Cardiac and Arterial Vascular Occlusive Events: Evaluate cardiovascular status, monitor and manage cardiovascular risk factors during nilotinib therapy. (5.4)
- Pancreatitis and Elevated Serum Lipase: Monitor serum lipase; if elevations are accompanied by abdominal symptoms, interrupt doses and consider appropriate diagnostics to exclude pancreatitis. (5.5)
- Hepatotoxicity: Monitor hepatic function tests monthly or as clinically indicated. (5.6)
- Electrolyte Abnormalities: Monitor nilotinib can cause hypophosphatemia, hypokalemia, hypocalcemia, and hyponatremia. Correct electrolyte abnormalities prior to initiating nilotinib and monitor periodically during therapy. (5.7)
- Tumor Lysis Syndrome: Maintain adequate hydration and correct uric acid levels prior to initiating therapy with nilotinib. (5.8)
- Hemorrhage: Hemorrhage from any site may occur. Advise patients to report signs and symptoms of bleeding and medically manage as needed. (5.9)
- Fluid Retention: Monitor patients for unexpected rapid weight gain, swelling, and shortness of breath. Manage medically. (5.13)
- Effects on Growth and Development in Pediatric Patients: Growth retardation has been reported in pediatric patients treated with nilotinib. Monitor growth and development in pediatric patients. (5.14)
- Embryo-Fetal Toxicity: Advise females of reproductive potential of potential risk to a fetus and to use effective contraception. (5.15, 8.1, 8.3)
- Treatment Discontinuation: Patients must have typical BCR-ABL transcripts. An FDA-authorized test with a detection limit below MR4.5 must be used to determine eligibility for discontinuation. Patients must be frequently monitored by the FDA authorized test to detect possible loss of remission. (5.16)

ADVERSE REACTIONS

The most common non-hematologic adverse reactions ($\geq 20\%$) in adult patients were nausea, rash, headache, fatigue, pruritus, vomiting, diarrhea, cough, constipation, arthralgia, nasopharyngitis, pyrexia, and night sweats. Hematologic adverse reactions included myelosuppression (thrombocytopenia, neutropenia, and anemia). (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Helsir Labs Limited at 1-866-495-1995 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Strong CYP3A4 Inhibitors: Avoid concomitant use with nilotinib, or reduce nilotinib dose if coadministration cannot be avoided. (7.1)
- Strong CYP3A4 Inducers: Avoid concomitant use with nilotinib. (7.1)
- Proton Pump Inhibitors: Use short-acting antacids or H2 blockers as an alternative to proton pump inhibitors. (7.1)

USE IN SPECIFIC POPULATIONS

- Lactation: Advise women not to breastfeed. (8.2)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Additional pediatric use information is approved for Novartis Pharmaceuticals Corporation's TASIGNA (nilotinib) capsules. However, due to Novartis Pharmaceuticals Corporation's marketing exclusivity rights, this drug product is not labeled with that pediatric information.

Revised: 05/2024

6.2	Postmarketing Experience
7	DRUG INTERACTIONS
7.1	Effect of Other Drugs on Nilotinib
7.2	Drugs That Prolong the QT Interval
8	USE IN SPECIFIC POPULATIONS
8.1	Pregnancy
8.2	Lactation
8.3	Females and Males of Reproductive Potential
8.4	Pediatric Use
8.5	Geriatric Use
8.6	Cardiac Disorders
8.7	Hepatic Impairment
10	OVERDOSAGE
11	DESCRIPTION
12	CLINICAL PHARMACOLOGY
12.1	Mechanism of Action
12.2	Pharmacodynamics
12.3	Pharmacokinetics
12.5	Pharmacogenomics
13	NONCLINICAL TOXICOLOGY
13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility
14	CLINICAL STUDIES
14.1	Adult Newly Diagnosed Ph+ CML-CP
14.2	Adult Patients With Resistant or Intolerant Ph+ CML-CP and CML-AP
14.3	Treatment Discontinuation in Newly Diagnosed Ph+ CML-CP Patients Who Have Achieved a Sustained Molecular Response (MR4.5)
14.4	Treatment Discontinuation in Ph+ CML-CP Patients Who Have Achieved a Sustained Molecular Response (MR4.5) on Nilotinib Following Prior Imatinib Therapy
16	HOW SUPPLIED/STORAGE AND HANDLING
17	PATIENT COUNSELING INFORMATION
* Sections or subsections omitted from the full prescribing information are not listed.	

