



Manufactured for:
Ascent Pharmaceuticals, Inc.
Central Islip, NY 11722

Manufactured for:
Camber Pharmaceuticals, Inc.
Piscataway, NJ 08854

Rev: 01/2026



Manufactured by:
Ascent Pharmaceuticals, Inc.
Central Islip, NY 11722



NDC 31722-482-30

Buprenorphine and Naloxone Sublingual Film



12 mg/3 mg per Sublingual Film

Children who accidentally take buprenorphine and naloxone sublingual film will need emergency medical care.

Keep buprenorphine and naloxone sublingual film out of the reach of children.

Do not cut, chew or swallow sublingual film.

Instructions for the Pharmacist: Dispense the enclosed Medication Guide to each patient.

30 pouches each containing 1 sublingual film

Rx Only

NO VARNISH

Buprenorphine and Naloxone
Sublingual Film
12 mg/3 mg per Sublingual Film

Nick

Buprenorphine and Naloxone Sublingual Film



12 mg/3 mg per Sublingual Film

Each Sublingual Film contains buprenorphine 12 mg and naloxone 3 mg

Taking buprenorphine and naloxone sublingual film

- If directed to use 2 films, place films on opposite sides under the tongue or inside the cheek. Avoid letting films touch
- Keep in place until fully dissolved
- If told to use a third film, use as directed after the first 2 films have dissolved
- See package insert for dosing information
- Always take as prescribed

Sublingual Administration:

- Place film under tongue (close to the base on right or left side)

Buccal Administration:

- Place film on inside of right or left cheek

Store at 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

Children who accidentally take buprenorphine and naloxone sublingual film will need emergency medical care. Keep buprenorphine and naloxone sublingual film out of the reach of children.

Medication Guide available at
<http://camberpharma.com/medication-guides>

LEBG3043C2
center seam